


(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
13 November 2003 (13.11.2003)

PCT

(10) International Publication Number
WO 03/092600 A2

- (51) International Patent Classification⁷: **A61K** (74) Agent: **FIELD, Bret E.**; Bozicevic, Field & Francis, 200 Middlefield Road, Suite 200, Menlo Park, CA 94025 (US).
- (21) International Application Number: PCT/US03/13492 ✓ (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 29 April 2003 (29.04.2003) ✓
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 10/136,860 30 April 2002 (30.04.2002) US ✓ (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; 1111 Franklin Street, 12th Floor, Oakland, CA 94607-5200 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **PORTNOY, Daniel** ✓ A. [US/US]; 1196 Curtis Street, Albany, CA 94706 (US). **CALENDAR, Richard** [US/US]; 940 Euclid Avenue, Berkeley, CA 94708 (US). **LAUER, Peter M.** [US/US]; 5719 Oak Grove Avenue, Oakland, CA 94618 (US).
- Published: — without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SITE SPECIFIC LISTERIA INTEGRATION VECTORS AND METHODS FOR USING THE SAME ✓



(57) Abstract: Site-specific *Listeria* integration vectors and methods for their use are provided. The subject vectors include a bacteriophage integrase gene and a bacteriophage attachment site, where in many embodiments the bacteriophage that is the source of these elements is a listeriophage. In certain embodiments, the subject vectors further include a multiple cloning site, where the multiple cloning site may further include a polypeptide coding sequence, e.g., for a heterologous antigen. The subject vectors and methods find use in a variety of different applications, including the study of *Listeria* species and the preparation of *Listeria* vaccines.

WO 03/092600 A2

4/pet3

SITE SPECIFIC LISTERIA INTEGRATION VECTORS AND METHODS FOR USING THE SAME

ACKNOWLEDGMENT OF GOVERNMENT SUPPORT

5

This invention was made with Government support under Grant Nos. 1 R37 AI29619 and 1 R01 AI27655 awarded by the National Institute of Health. The Government has certain rights in this invention.

10

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of application serial no. 10/136,860 filed April 30, 2002; the disclosure of which is herein incorporated by reference.

15

INTRODUCTION

Field of the Invention

The field of this invention is *Listeria* species, e.g., *Listeria monocytogenes*, particularly recombinant strains of *Listeria* species, and methods for their
20 fabrication and use.

Background of the Invention

Listeria monocytogenes is a Gram-positive, food-borne human and animal pathogen responsible for serious infections in immunocompromised individuals and pregnant women. Severe *L. monocytogenes* infections in humans are
25 characterized by meningitis, meningoencephalitis, septicemia, and fetal death. *L. monocytogenes* is ubiquitous in nature and, in addition, can be isolated from a wide variety of warm blooded animals.

Protocols for recombinantly engineering *Listeria* species are of interest in both research and therapeutic applications. For example, *Listeria* species
30 transformation protocols find use in the elucidation of the mechanisms responsible for growth and virulence of these types of bacteria. In addition, such protocols also find use in the preparation of live *Listeria* vaccines, which vaccines find use in a variety of different medical applications.

To date, a variety of different protocols have been employed to transform *Listeria* species, where such protocols include: homologous recombination, transposon based recombination, etc. While different protocols are currently available for engineering *Listeria* species, such methods are not entirely
5 satisfactory. Disadvantages currently experienced with one or more of the available protocols include: (1) instability of the expression cassette in the transformed species; (2) variable impact on virulence of the transformed species; (3) size constraints of the expression cassette that can be placed in the transformed species; etc.

10 As such, despite the variety of different transformation protocols available, there is continued interest in the identification of further transformation protocols that can expand the repertoire of available genetic tools. Of particular interest would be the development of an efficient, site-specific integration vector that was suitable for use with a wide array of different *Listeria* species, where the vector
15 did not suffer from one or more of the above disadvantages of the currently available protocols.

Relevant Literature

Patents and published patent applications of interest include: U.S. Patent No. 5,830,702 and published PCT application serial nos.: WO 99/25376 and WO
20 00/09733.

SUMMARY OF THE INVENTION

Site-specific *Listeria* species integration vectors and methods for their use are provided. The subject vectors include a bacteriophage integrase gene and a bacteriophage attachment site, where in many embodiments the bacteriophage
25 that is source of these elements is a listeriphage. In certain embodiments, the subject vectors further include a multiple cloning site, where the multiple cloning site may further include a coding sequence, e.g., a coding sequence for a heterologous polypeptide, etc. The subject vectors and methods find use in a variety of different applications, including the study of *Listeria* species and the
30 preparation of *Listeria* species. vaccines.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1. (A) Plasmid map of the pPL1 integration vector. SEQ ID NO:24 provides the sequence of pPL1. Chloramphenicol resistance genes and *E. coli*

origin of replication are shown in grey, RP4 origin of transfer shown in white, integrase gene and *L. monocytogenes* p60 promoter shown in black. Multiple cloning site is shown at the bottom of the plasmid with unique restriction sites noted below the MCS in a box. pPL24 and pPL25 inserts are shown

5 schematically below the MCS and were cloned as described in Materials and Methods. Final sizes of the plasmid constructs and the restriction sites used in cloning are noted with each of the inserts. (B) Plasmid map of the integration vector pPL2. SEQ ID NO:28 provides the sequence of pPL2. The color scheme and genes are the same as in FIG. 1A except the PSA integrase and PSAattPP' sites as noted. The multiple cloning site with 13 unique restriction sites is shown
10 at the bottom of the plasmid.

FIG. 2. Genomic organization of the attachment sites within the *comK* gene (A and B) and the tRNA^{Arg} gene (C and D). (A) Non-lysogenic *L. monocytogenes* strain, with an intact *comK* gene. Primers PL60 and PL61
15 amplify across the bacterial attachment site *comK-attBB'*. (B) Lysogenic *L. monocytogenes* strain (with approximately 40 kb of phage DNA inserted into the *comK* gene) or integrated strain (with pPL1 construct inserted into the *comK* gene). Primers PL14 and PL61 amplify across the hybrid attachment site *comK-attPB'*. (C) *L. monocytogenes* serotype 4b strain non-lysogenic at tRNA^{Arg}-*attBB'*.
20 Primers NC16 and NC17 amplify across the bacterial attachment site tRNA^{Arg}-*attBB'* in serotype 4b strains. Asterisk indicates primers NC16 and NC17 are substituted with PL102 and PL103 to amplify across the bacterial attachment site tRNA^{Arg}-*attBB'* in serotype 1/2 strains. (D) Lysogenic *L. monocytogenes* strain (with approximately 40 kb of phage DNA or 6 kb pPL2 vector DNA inserted at the
25 3' end of the tRNA^{Arg} gene. Primers NC16 and PL95 amplify across the hybrid attachment site tRNA^{Arg}-*attBP'* in both serotype 4b and 1/2 strains.

FIG. 3. Expression and functional complementation of ActA in SLCC-5764. (A) Coomassie blue stained SDS-PAGE of SLCC-5764 derived strains grown to late-log phase. ActA is indicated by an arrow. Lane 1: molecular size
30 marker; lane 2: DP-L3780; lane 3: DP-L4083; lane 4: DP-L4086; lane 5: SLCC-5764; lane 6: DP-L4082; lane 7: DP-L4084; lane 8: DP-L4085; lane 9: DP-L4087. Strains are described in Table 1. (B) Actin tail formation and movement of DP-

L4087 in *Xenopus* cell extract. The top panel is a phase image; the bottom panel is a fluorescent image of the same field.

FIG 4. (A) Clover-leaf diagram of tRNA^{Arg} utilized as the PSA attachment site. The arginine anticodon is circled. The region with sequence identity between the tRNA gene and the PSA *attPP'* is outlined. The boundaries of the tRNA^{Arg} gene and Cove score (82.37) were predicted with tRNAscan-SE (31). (B) Alignment of the tRNA^{Arg}-*attBB'* region of *L. monocytogenes* WSLC 1042 (top line) and the *attPP'* region of PSA downstream of the integrase gene. The 74 nt tRNA^{Arg} gene of *L. monocytogenes* is boxed and the 17 bp overlapping region of sequence identity (core integration site) is shaded grey. The tRNA^{Arg} gene anticodon is shown in bold and underlined. Identical nucleotide residues are indicated by (:). The numbers located at the left indicate the nucleotide position in the DNA sequences of the WSLC 1042 attachment site (AJ314913) and PSA genome (AJ312240).

DESCRIPTION OF THE SPECIFIC EMBODIMENTS

Site-specific *Listeria* species integration vectors and methods for their use are provided. The subject vectors include a bacteriophage integrase gene and a bacteriophage attachment site, where in many embodiments the bacteriophage that is source of these elements is a listeriophage. In certain embodiments, the subject vectors further include a multiple cloning site, where the multiple cloning site may further include a coding sequence, e.g., for a heterologous polypeptide, etc. The subject vectors and methods find use in a variety of different applications, including the study of *Listeria* species and the preparation of *Listeria* species vaccines.

Before the subject invention is described further, it is to be understood that the invention is not limited to the particular embodiments of the invention described below, as variations of the particular embodiments may be made and still fall within the scope of the appended claims. It is also to be understood that the terminology employed is for the purpose of describing particular embodiments, and is not intended to be limiting. Instead, the scope of the present invention will be established by the appended claims.

In this specification and the appended claims, the singular forms "a," "an" and "the" include plural reference unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs.

Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range, and any other stated or intervening value in that stated range, is encompassed within the invention.

The upper and lower limits of these smaller ranges may independently be included in the smaller ranges, and are also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in the invention.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs. Although any methods, devices and materials similar or equivalent to those described herein can be used in the practice or testing of the invention, representative methods, devices and materials are now described.

All publications mentioned herein are incorporated herein by reference for the purpose of describing various invention components that are described in the publications which might be used in connection with the presently described invention.

In further describing the subject invention, the subject vectors are reviewed first in greater detail, followed by a review of the methods of using the subject vectors, as well as representative applications in which the subject vectors and methods find use.

30

VECTORS

As summarized above, the subject invention provides *Listeria* site-specific integration vectors, i.e., vectors that integrate into *Listeria* genomes in a site-

specific manner. The subject vectors are characterized in that they stably integrate into a genome of a *Listeria* species in a site-specific, as opposed to random, manner. As the subject vectors integrate site-specifically into a target genome, the vectors of the subject invention typically are inserted into a specific location or sequence (i.e., domain, location) of the target genome by a mechanism mediated by a recombinase, specifically an integrase, where the integrase is one that uses two distinct recognition sites as substrates, one of which is positioned in the integration site of the target genome (the site into which a nucleic acid is to be integrated) and another adjacent a nucleic acid of interest to be introduced into the integration site (i.e., a site on the subject vector referred to herein as the "bacteriophage attachment site"). For example, the recognition sites for phage integrases are generically referred to as attB, which is present in the bacterial genome (into which nucleic acid is to be inserted) and attP (which is present in the phage nucleic acid adjacent the nucleic acid for insertion into the bacterial genome, which is referred to herein as the "bacteriophage attachment site"). Recognition sites can be native (endogenous to a target genome) or altered relative to a native sequence. Use of the term "recognize" in the context of an integrase "recognizes" a recognition sequence, is meant to refer to the ability of the integrase to interact with the recognition site and facilitate site-specific recombination.

The subject vectors are capable of integrating into the genomes of a wide variety of different *Listeria* species. Integration is readily determined by using any convenient assay, including those known to those of skill in the art, that can identify whether or not vector DNA has integrated into the genomic DNA of a target organism. For example, vector DNA can be introduced into a target organism, e.g., via conjugation, and then integration can be determined via PCR amplification of the integrated sequence using primers flanking the target site, where the size of the amplification product is determinative of whether integration has occurred. An example of such an assay is provided in the experimental section, below. Additional features of many embodiments of the subject vectors is that they replicate autonomously in a non-*Listeria* host cell, e.g., *E. coli*, and are stable and innocuous in such non-*Listeria* host cells.

The subject integration vectors are typically double-stranded plasmid vectors, where the vectors are generally at least about 3 kb, often at least about 5

kb and may be large as 15 kb, 20 kb, 25 kb, 30 kb or larger, where the vectors sometimes range in size from about 3-6 to about 20 kb. The vectors include a number of structural features that impart to the vectors the above-described functional characteristics.

5 One structural feature of the subject vectors is a bacteriophage integrase gene, i.e., a nucleic acid coding sequence for a bacteriophage integrase, which is operably linked to a *Listeria* specific promoter, such that the gene (coding sequence) is expressed in the *Listeria* cell for which the vector is designed to be employed. In many embodiments, the bacteriophage integrase gene is one
10 obtained from a listeriophage, i.e., it is a listeriophage integrase gene. A variety of different listeriophages are known in the art, where any convenient listeriophage may serve as the source of the integrase gene, i.e., the integrase encoding nucleic acid. Specific integrases of interest include, but are not limited to: the U153 integrase, the PSA integrase; and the like.

15 As indicated above, the integrase gene is operably linked to a promoter (as well as regulatory and/or signal sequences, if necessary) that drives expression of the integrase gene when the vector is present in the *Listeria* cell for which the vector is designed and in which integration of the vector is desired. Any convenient promoter may be employed, where in certain embodiments the
20 promoter is a *Listeria* specific promoter. Representative promoters of interest include, but are not limited to: the *Listeria* p60 promoter, the *Listeria actA* promoter, the *Listeria plcA* promoter, the *Listeria mpl* promoter, the *Listeria plcB* promoter, the *Listeria inlA* promoter; a heat shock promoter; and the like. Promoters may also include certain bacteriophage promoters such as the
25 promoters for T7, Q β , SP6 and the like if the strain of *Listeria* also expresses the cognate bacteriophage RNA polymerase. In addition to the integrase gene/promoter element described above, the subject vectors also include a phage attachment site, i.e., a sequence or domain of nucleotides that provide for a site-specific integration with a *Listeria* genome. Any convenient phage
30 attachment site may be employed, where selection of the phage attachment site will depend, at least in part, on the desired integration location for the vector. Representative phage attachment sites of interest include, but are not limited to: the comK integration site (as described in greater detail in the experimental

section below); the tRNA^{Arg} integration site (as described in greater detail in the experimental section below); and the like.

In addition, the subject vectors may include an origin of replication that provides for replication of the vector in a non-*Listeria* host cell, e.g., *E.coli*. This
5 origin of replication may be any convenient origin of replication or ori site, where a number of ori sites are known in the art, where particular sites of interest include, but are not limited to: p15A; pSC101; ColEI; pUC; pMB9; and the like.

In addition, the subject vectors may include an origin of transfer site or element when convenient, e.g., when the vector is introduced in the target
10 *Listeria* cell using a conjugation protocol, as described in greater detail below. Any convenient origin of transfer (oriT) may be employed, where representative origins of transfer of interest include, but are not limited to: RP4 oriT; RSF1010 oriT; and the like.

15 In addition, the subject vectors typically include at least one restriction endonuclease recognized site, i.e., restriction site, which is located on the vector at a location which is amenable to insertion of a heterologous gene/expression cassette. A variety of restriction sites are known in the art and may be present on the vector, where such sites include those recognized by the following restriction
20 enzymes: *HindIII*, *PstI*, *Sall*, *AccI*, *HincII*, *XbaI*, *BamHI*, *SmaI*, *XmaI*, *KpnI*, *SacI*, *EcoRI*, *BstXI*, *EagI*, *NotI*, *SpeI* and the like. In many embodiments, the vector includes a polylinker or multiple cloning site, i.e., a closely arranged series or array of sites recognized by a plurality of different restriction enzymes, such as those listed above.

25 In certain embodiments, the subject vectors include at least one coding sequence, e.g., a coding sequence for heterologous polypeptide/protein coding sequence present in the multiple cloning site, e.g., as a result of using a restriction endonuclease site present in the multiple cloning site to insert the coding sequence into the vector, according to well known recombinant
30 technology protocols. By "heterologous" is meant that the coding sequence encodes a product, e.g., a protein, peptide, polypeptide, glycoprotein, lipoprotein, or other macromolecule, that is not normally expressed in *Listeria*. In many embodiments, this coding sequence is part of an expression cassette, which provides for expression of the coding sequence in the *Listeria* cell for which the

vector is designed. The term "expression cassette" as used herein refers to an expression module or expression construct made up of a recombinant DNA molecule containing at least one desired coding sequence and appropriate nucleic acid sequences necessary for the expression of the operably linked
5 coding sequence in a particular host organism, i.e., the *Listeria* cell for which the vector is designed, such as the promoter/regulatory/signal sequences identified above, where the expression cassette may include coding sequences for two or more different polypeptides, or multiple copies of the same coding sequence, as desired. The size of the coding sequence and/or expression cassette that
10 includes the same may vary, but typically falls within the range of about 25-30 to about 6000 bp, usually from about 50 to about 2000 bp. As such, the size of the encoded product may vary greatly, and a broad spectrum of different products may be encoded by the expression cassettes present in the vectors of this embodiment.

15 The nature of the coding sequence and other elements of the expression cassette may vary, depending on the particular application of the vector, e.g., to study *Listeria* species, to produce *Listeria* species vaccines, for cytosolic delivery of macromolecules, etc. For example, where the vectors are employed in the production of *Listeria* vaccines, the coding sequence may encode a heterologous
20 antigen, where representative heterologous antigens of interest include, but are not limited to: (a) viral antigens, e.g., influenza np protein, HIV gag protein, HIV env protein or parts thereof, e.g., gp120 and gp41, HIV nef protein, HIV pol proteins, HIV reverse transcriptase, HIV protease, herpes virus proteins, etc., (b) malarial antigens; (c) fungal antigens; (d) bacterial antigens; (e) tumor and tumor
25 related antigens; and the like. Due to the flexibility of the vector system, virtually any coding sequence of interest may be inserted. Where secretion of the product encoded by the expression cassette is desired, the expression cassette may include a coding sequence for a fusion protein of a selected foreign antigen and a protein that directs secretion, e.g., Listeriolysin O or PI-PLC; a signal sequence,
30 such as hemolysin signal sequence, etc. Where the subject vectors are employed in the preparation of *Listeria* delivery vehicles, e.g., as described in PCT publication no. WO 00/09733 (the priority application of which is herein incorporated by reference); and Dietrich et al., *Nature Biotechnology* (1998) 16: 181-185, the heterologous polypeptide coding sequence may be a cytolysin, e.g.,

phospholipase, pore forming toxin, listeriolysin O, streptolysin O, perfringolysin O, acid activated cytolysins, phage lysins, etc. Other coding sequences of interest include, but are not limited to: cytokines, costimulatory molecules, and the like. As indicated above, the vector may include at least one coding sequence, where
5 in certain embodiments the vectors include two or more coding sequences, where the coding sequences may encode products that act concurrently to provide a desired results.

In general, the coding sequence may encode any of a number of different products and may be of a variety of different sizes, where the above discussion
10 merely provides representative coding sequences of interest.

METHODS OF PREPARING THE SUBJECT VECTORS

The vectors of the subject invention may be produced using any
15 convenient protocol, including by standard methods of restriction enzyme cleavage, ligation and molecular cloning. One protocol for constructing the subject vectors includes the following steps. First, purified nucleic acid fragments containing desired component nucleotide sequences as well as extraneous sequences are cleaved with restriction endonucleases from initial sources.
20 Fragments containing the desired nucleotide sequences are then separated from unwanted fragments of different size using conventional separation methods, e.g., by agarose gel electrophoresis. The desired fragments are excised from the gel and ligated together in the appropriate configuration so that a circular nucleic acid or plasmid containing the desired sequences, e.g. sequences corresponding
25 to the various elements of the subject vectors, as described above is produced. Where desired, the circular molecules so constructed are then amplified in a host, e.g. *E. coli*. The procedures of cleavage, plasmid construction, cell transformation and plasmid production involved in these steps are well known to one skilled in the art and the enzymes required for restriction and ligation are
30 available commercially. (See, for example, R. Wu, Ed., Methods in Enzymology, Vol. 68, Academic Press, N.Y. (1979); T. Maniatis, E. F. Fritsch and J. Sambrook, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1982); Catalog 1982-83, New England Biolabs, Inc.; Catalog 1982-83, Bethesda Research Laboratories, Inc. An example of how to

construct a vector of the present invention is provided in the Experimental section, below.

METHODS

5

Also provided by the subject invention are methods of using the above described *Listeria* site specific integration vectors to integrate a heterologous nucleic acid into a *Listeria* genome. In practicing the subject methods, a vector of the subject invention is introduced into a target *Listeria* cell under conditions
10 sufficient for integration of the vector into the target cell genome to occur. Any convenient protocol for introducing the vector into the target cell may be employed.

Suitable protocols include: calcium phosphate mediated transfection; protoplast fusion, in which protoplasts harboring amplified amounts of vector are
15 fused with the target cell; electroporation, in which a brief high voltage electric pulse is applied to the target cell to render the cell membrane of the target cell permeable to the vector; microinjection, in which the vector is injected directly into the cell, as described in Capechhi et al, *Cell* (1980) 22: 479; and the like. The above *in vitro* protocols are well known in the art and are reviewed in greater
20 detail in Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press)(1989) pp16.30-16.55.

In certain embodiments, e.g., where direct introduction into the target *Listeria* cell does not provide optimal results, one representative method that may be employed is a conjugation method, which comprises: (a) introducing the
25 vector into a non-*Listeria* host cell, e.g., *E. coli*; followed by (b) transfer of the vector from the non-*Listeria* host cell to the *Listeria* host cell, e.g., by conjugation. Introduction into the non-*Listeria* host cell may be accomplished using any of the protocols described above. For the conjugation step, any convenient protocol may be employed, where suitable protocols typically include combining the donor
30 and acceptor cells in a suitable medium and maintaining under suitable conditions for conjugation and vector transfer to occur. Specific representative conditions are provided in the Experimental section below.

The above methods result in stable integration of the vector and any expression cassette carried thereby, e.g., that encodes a heterologous

protein/foreign antigen, into a target *Listeria* cell genome in a site specific manner. The subject methods find use with a wide variety of different *Listeria* species.

5 UTILITY

The above described vectors and methods of using the same find use in a variety of different applications, where representative applications include, but are not limited to: (a) research applications; (b) vaccine preparation applications; (c) 10 *Listerial* delivery vehicle preparation applications; and the like.

One type of application in which the subject vectors and methods of using may be employed is in research of *Listeria* species. For example, the subject vectors and methods allow simple and efficient strain construction and are widely useful in various strains used to study the intracellular life cycle of *L. monocytogenes*. 15 Additionally, the subject vectors and methods may be employed to produce stable merodiploid strains to allow refined copy number studies and studies of interactions within a protein through multimerization and testing of the dominance or recessive nature of different alleles of a gene in the same bacterial strain.

20 The subject vectors and methods also find use in vaccine preparation applications. For example, the subject vectors find use in the production of *Listeria* cultures capable of expressing a heterologous antigen, i.e., *Listeria* vaccines, where the *Listeria* cells employed may be attenuated. The attenuated strains employed may be capable of normal invasion of a host cell, but incapable 25 of normal survival or growth in the cell or cell-to-cell spread, or they may have other alterations that preclude normal pathogenicity.

The vaccines produced using vectors of the present invention are administered to a vertebrate by contacting the vertebrate with a sublethal dose of the genetically engineered *Listeria* vaccine, where contact typically includes 30 administering the vaccine to the host. Thus the present invention provides for vaccines genetically engineered with the integration vector and provided in a pharmaceutically acceptable formulation. Administration can be oral, parenteral, intranasal, intramuscular, intravascular, direct vaccination of lymph nodes, administration by catheter or any one or more of a variety of well-known

administration routes. In farm animals, for example, the vaccine may be administered orally by incorporation of the vaccine in feed or liquid (such as water). It may be supplied as a lyophilized powder, as a frozen formulation or as a component of a capsule, or any other convenient, pharmaceutically acceptable formulation that preserves the antigenicity of the vaccine. Any one of a number of well known pharmaceutically acceptable diluents or excipients may be employed in the vaccines of the invention. Suitable diluents include, for example, sterile, distilled water, saline, phosphate buffered solution, and the like. The amount of the diluent may vary widely, as those skilled in the art will recognize. Suitable excipients are also well known to those skilled in the art and may be selected, for example, from A. Wade and P.J. Weller, eds., *Handbook of Pharmaceutical Excipients* (1994) The Pharmaceutical Press: London. The dosage administered may be dependent upon the age, health and weight of the patient, the type of patient, and the existence of concurrent treatment, if any. The vaccines can be employed in dosage forms such as capsules, liquid solutions, suspensions, or elixirs, for oral administration, or sterile liquid for formulations such as solutions or suspensions for parenteral, intranasal intramuscular, or intravascular use. In accordance with the invention, the vaccine may be employed, in combination with a pharmaceutically acceptable diluent, as a vaccine composition, useful in immunizing a patient against infection from a selected organism or virus or with respect to a tumor, etc. Immunizing a patient means providing the patient with at least some degree of therapeutic or prophylactic immunity against selected pathogens, cancerous cells, etc.

The subject vaccines prepared with the subject vectors find use in methods for eliciting or boosting a helper T cell or a cytotoxic T-cell response to a selected agent, e.g., pathogenic organism, tumor, etc., in a vertebrate, where such methods include administering an effective amount of the Listeria vaccine. The subject vaccines prepared with the subject vectors find use in methods for eliciting in a vertebrate an innate immune response that augments the antigen-specific immune response. Furthermore, the vaccines of the present invention may be used for treatment post-exposure or post diagnosis. In general, the use of vaccines for post-exposure treatment would be recognized by one skilled in the art, for example, in the treatment of rabies and tetanus. The same vaccine of the present invention may be used, for example, both for immunization and to boost

immunity after exposure. Alternatively, a different vaccine of the present invention may be used for post-exposure treatment, for example, such as one that is specific for antigens expressed in later stages of exposure. As such, the subject vaccines prepared with the subject vectors find use as both prophylactic and
5 therapeutic vaccines to induce immune responses that are specific for antigens that are relevant to various disease conditions.

The patient may be any human and non-human animal susceptible to infection with the selected organism. The subject vaccines will find particular use with vertebrates such as man, and with domestic animals. Domestic animals
10 include domestic fowl, bovine, porcine, ovine, equine, caprine, Leporidate (such as rabbits), or other animal which may be held in captivity.

In general, the subject vectors and methods find use in the production of vaccines as described U.S. Patent No. 5,830,702, the disclosure of which is herein incorporated by reference; as well as PCT publication no W0 99/25376,
15 the disclosures of the priority applications of which are herein incorporated by reference.

The subject vectors also find use in the production of listerial delivery vehicles for delivery of macromolecules to target cells, e.g., as described in: PCT publication no. WO 00/09733 (the priority application of which is herein
20 incorporated by reference); and Dietrich et al., Nature Biotechnology (1998) 16: 181-185. A variety of different types of macromolecules may be delivered, including, but not limited to: nucleic acids, polypeptides/proteins, etc., as described in these publications.

KITS & SYSTEMS

25 Also provided are kits and systems that find use in preparing the subject vectors and/or preparing recombinant Listeria cells using the subject vectors and methods, as described above. For example, kits and systems for producing the subject vectors may include one or more of: an initial vector with a multiple
30 cloning site; a restriction endonuclease for cleaving a site in the multiple cloning site, a vector including an expression cassette of interest which is to be inserted into the multiple cloning site; etc. Where the kits and systems are designed for the production of the recombinant Listeria cells, the kits and systems may include

vectors, or components for making the same, as described above, *Listeria* target cells, non-*Listeria* host cells, and the like. The subject kits may further include other components that find use in the subject methods, e.g., reaction buffers, growth mediums, etc.

- 5 The various reagent components of the kits may be present in separate containers, or may all be precombined into a reagent mixture for combination with template DNA.

 In addition to the above components, the subject kits will further include instructions for practicing the subject methods. These instructions may be
10 present in the subject kits in a variety of forms, one or more of which may be present in the kit. One form in which these instructions may be present is as printed information on a suitable medium or substrate, e.g., a piece or pieces of paper on which the information is printed, in the packaging of the kit, in a package insert, etc. Yet another means would be a computer readable medium,
15 e.g., diskette, CD, etc., on which the information has been recorded. Yet another means that may be present is a website address which may be used via the internet to access the information at a removed site. Any convenient means may be present in the kits.

- 20 The following examples are offered by way of illustration and not by way of limitation.

EXPERIMENTAL

25 I. MATERIALS AND METHODS

A. Construction of pPL1 integration vector.

Standard molecular techniques were used in the construction of the 6101 bp integration vector pPL1 (Fig. 1A). The complete sequence of the pPL1 vector
30 is provided as SEQ ID NO:25. pPL1 is a low copy plasmid that replicates autonomously in *E. coli* and integrates in a site-specific manner in *L. monocytogenes*, and was assembled from 6 independent DNA sources as follows. Restriction sites in the PCR primers used for construction are

underlined. All PCR reactions used in cloning steps utilized Vent DNA polymerase (New England Biolabs).

- The multiple cloning site (MCS) from pBluescript KS- (Alting-Mees, M. A., and J. M. Short. 1989. pBluescript II: gene mapping vectors. Nucleic Acids Res. 17:9494.) (bp 1-171) was cloned after PCR with primers 5'-GGACGTCATTAACCCTCACTAAAGG-3' and 5'-GGACGTCAATACGACTCACTATAGG-3' (SEQ ID NOS: 01 & 02). The low copy Gram-negative origin of replication and chloramphenicol acetyltransferase (CAT) gene from pACYC184 (Chang, A. C., and S. N. Cohen. 1978. Construction and characterization of amplifiable multicopy DNA cloning vehicles derived from the P15A cryptic miniplasmid. J. Bacteriol. 134:1141-1156.) (bp 172-2253) were cloned after PCR with primers 5'-GGACGTCGCTATTTAACGACCCTGC-3' and 5'-GAGCTGCAGGAGAATTACAACCTATATCGTATGGGG-3' (SEQ ID NOS: 03 & 04). For direct conjugation from *E. coli* to *L. monocytogenes*, the RP4 origin of transfer (*oriT*) (Pansegrau, W., E. Lanka, P. T. Barth, D. H. Figurski, D. G. Guiney, D. Haas, D. R. Helinski, H. Schwab, V. A. Stanisich, and C. M. Thomas. 1994. Complete nucleotide sequence of Birmingham IncP alpha plasmids. Compilation and comparative analysis. J. Mol. Biol. 239:623-663) (bp 2254-2624) was cloned from plasmid pCTC3 (Williams, D. R., D. I. Young, and M. Young. 1990. Conjugative plasmid transfer from *Escherichia coli* to *Clostridium acetobutylicum*. J. Gen. Microbiol. 136:819-826) after PCR with primers 5'-GCACTGCAGCCGCTTGCCCTCATCTGTTACGCC-3' and 5'-CATGCATGCCTCTCGCCTGTCCCCTCAGTTCAG-3' (SEQ ID NOS: 05 & 06). The listeriphage U153 integrase gene and attachment site (*attPP*) (A. Nolte, P. Lauer, and R. Calendar, unpublished; bp 2629-4127)(SEQ ID NO: 25 includes the sequences for these sites), that direct the site-specific integration of the plasmid were cloned after PCR with primers 5'-GTAGATCTTAACCTTCCATGCGAGAGGAG-3' and 5'-GGGCATGCGATAAAAAGCAATCTATAGAAAAACAGG-3' (SEQ ID NOS: 07 & 08). For expression of the U153 integrase gene, the *L. monocytogenes* p60 promoter (Lauer, P., J. A. Theriot, J. Skoble, M. D. Welch, and D. A. Portnoy. 2001. Systematic mutational analysis of the amino-terminal domain of the *Listeria monocytogenes* ActA protein reveals novel functions in actin-based motility. Mol. Microbiol. 42:1163-1177.) was PCR amplified with primers 5'-

CCTAAGCTTTTCGATCATCATAATTCTGTC-3' and 5'-

GGGCATGCAGATCTTTTTTTCAGAAAATCCCAGTACG-3' (SEQ ID NOS: 09 &

10) and cloned upstream of the integrase gene. Base pairs 4570-6101 are a

Hind III-*Aat* II restriction fragment subcloned from pUC18-Cat (a kind gift from

5 Nancy Freitag), and contain the inducible Gram-positive CAT gene from pC194

(Horinouchi, S., and B. Weisblum. 1982. Nucleotide sequence and functional

map of pC194, a plasmid that specifies inducible chloramphenicol resistance. J.

Bacteriol. 150:815-825) (bp 4788-5850).

10 B. Cloning of the *hly* and *actA* genes into pPL1.

The *hly* gene was subcloned from the plasmid pDP-906 (Jones, S., and D.

A. Portnoy. 1994. Characterization of *Listeria monocytogenes* pathogenesis in a

strain expressing perfringolysin O in place of listeriolysin O. Infect. Immun.

62(12):5608-5613.) by restriction digestion with *Bam*H I and *Xba* I, gel purifying a

15 2.9 kb fragment and by ligating it into pPL1 cut with *Bam*H I and *Spe* I. The

resultant plasmid was designated pPL24 (Fig. 1A). The *actA* gene was PCR

amplified from 10403S genomic DNA with primers 5'-

GGTCTAGATCAAGCACATACCTAG-3' and 5'-

CGGGATCCTGAAGCTTGGGAAGCAG-3' (SEQ ID NOS:11 & 12). The 2220 bp

20 PCR product was gel purified, cut with *Bam*H I and *Xba* I, and cloned into pPL1

cut with *Bam*H I and *Spe* I. The resultant plasmid was designated pPL25 (Fig.

1A).

25 C. Phage curing, conjugation and molecular confirmation of plasmid integration.

Phage curing was accomplished by adapting historical methodologies

(Cohen, D. 1959. A variant of Phage P2 Originating in *Escherichia coli*, strain B.

Virology 7:112-126; Six, E. 1960. Prophage substitution and curing in lysogenic

cells superinfected with hetero-immune phage. J. Bacteriol. 80:728-729.). *L.*

30 *monocytogenes* 10403S derivatives carrying a prophage at *comK-attBB'*

(integrated in the *comK* ORF as described (Loessner, M. J., R. B. Inman, P.

Lauer, and R. Calendar. 2000. Complete nucleotide sequence, molecular

analysis and genome structure of bacteriophage A118 of *Listeria*

monocytogenes: implications for phage evolution. Mol. Microbiol. 35(2):324-

340.)) were grown in BHI at 37°C to 10⁸ CFU/ml and infected with listeriphage U153 (Hodgson, D. A. 2000. Generalized transduction of serotype 1/2 and serotype 4b strains of *Listeria monocytogenes*. Mol. Microbiol. 35(2):312-323.) at a multiplicity of infection of 20:1 in the presence of 5 mM CaCl₂. Cultures were
5 incubated by shaking at 37°C for 75 minutes and inhibition of growth was monitored by comparison of the OD₆₀₀ of the infected culture with an uninfected control culture. The infected culture was diluted 10⁻² and 10⁻⁴ in BHI, and both dilutions were grown at 37°C until the 10⁻² dilution had increased 100-fold in optical density. The 10⁻⁴ fold dilution was then diluted 10⁻², and 3 µl were plated
10 on BHI. Fifty colonies were tested for phage release initially by toothpicking colonies into 0.25 ml LB broth and replica plating at 30°C on a lawn of Mack-4R (DP-L862), a non-lysogenic rough strain of *L. monocytogenes* particularly susceptible to forming plaques. Candidates that did not plaque were then tested by spotting 10 µl of culture on a lawn of Mack-4R to detect plaque formation. If
15 this second test was negative, the candidate was tested whether it could support plaque formation by the phage from the parent 10403S strain (Φ10403, (Hodgson, D. A. 2000. Generalized transduction of serotype 1/2 and serotype 4b strains of *Listeria monocytogenes*. Mol. Microbiol. 35(2):312-323.)). Curing was confirmed molecularly by PCR with the *comK-attBB'* specific primer pair
20 PL60/PL61 (sequences follow) for the absence of a phage at *comK-attBB'*. Approximately 10% of colonies were cured using this procedure.

Recipient strains of *L. monocytogenes* (SLCC-5764, DP-L1169 and DP-L1172) were made streptomycin resistant for counter-selection in conjugation experiments by plate selection on BHI supplemented with 200 µg/ml antibiotic.
25 pPL1 plasmid constructs were electroporated into *E. coli* strain SM10 (Simon, R., U. Priefer, and A. Pühler. 1983. A broad host range mobilization system for *in vitro* genetic engineering: transposon mutagenesis in Gram negative bacteria. Bio/Technology 1:784-791.) using standard techniques (Sambrook, J., T. Maniatis, and E. F. Fritsch. 1989. Molecular Cloning : A
30 Laboratory Manual, 2nd edn. ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.). Bacterial strains were grown to mid-log (OD₆₀₀ ~0.55) shaking at 30°C. *E. coli* donor strains were grown in LB containing 25 µg/ml of chloramphenicol, *L. monocytogenes* recipient strains were grown in BHI. 2.5 ml

of donor culture was mixed with 1.5 ml of recipient culture and filtered onto pre-washed 47 mm 0.45 µm HA type filters (Millipore). The filter was washed once with 10 ml BHI, transferred to a BHI plate with no antibiotics and incubated for 2 hours at 30°C. The bacterial cells were gently resuspended in 2.5 ml of BHI and 5 25 µl and 50 µl aliquots were plated in 3 ml of LB top agar on BHI plates supplemented with 7.5 µg/ml chloramphenicol and 200 µg/ml streptomycin. Plates were incubated at 30°C overnight and shifted to 37°C for 2-3 days. Individual colonies were picked and screened by PCR for integration at the phage attachment site using the primers PL14 (5'-CTCATGAACTAGAAAAATGTGG-3') 10 (SEQ ID NO:13), PL60 (5'-TGAAGTAAACCCGCACACGATG-3')(SEQ ID NO:14) and PL61 (5'-TGTAACATGGAGGTTCTGGCAATC-3')(SEQ ID NO:15). PCR reactions were performed on small portions of individual bacterial colonies picked with sterile P200 pipet tips from BHI plates directly into 20 µl PCR reactions. The primer pair PL14/PL61 specifically amplifies *attBP'* in a PCR reaction, resulting in 15 a 743 bp product on integrated strains (both prophage and pPL1 derivatives). The primer pair PL60/PL61 specifically amplifies *comK-attBB'* in a PCR reaction, resulting in a 417 bp product only on non-lysogenic strains (i.e. DP-L4056). PCR assays were performed in a Hybaid Omn-E thermocycler with an annealing temperature of 55°C for 30 cycles. Integrants arose at a frequency of 20 approximately 2×10^{-4} per donor cell.

D. Hemolysis on blood plates and hemolytic activity assay.

Hemolysis on blood plates was scored on tryptic soy agar plates supplemented with 5% defibrinated sheep blood (HemoStat, Davis CA).

25 Hemolytic assays were performed essentially as described (Portnoy, D. A., P. S. Jacks, and D. J. Hinrichs. 1988. Role of hemolysin for the intracellular growth of *Listeria monocytogenes*. J. Exp. Med. 167(4):1459-1471.). Hemolytic activity is expressed as the reciprocal of the dilution of culture supernatant required to lyse 50% of sheep erythrocytes.

30

E. Plaquing in L2 cells.

Plaque sizes were determined as previously described (Lauer, P., J. A. Theriot, J. Skoble, M. D. Welch, and D. A. Portnoy. 2001. Systematic mutational

analysis of the amino-terminal domain of the *Listeria monocytogenes* ActA protein reveals novel functions in actin-based motility. Mol. Microbiol. 42(5):1163-1177.). Each strain was plaqued in 6 to 8 independent experiments and compared to 10403S in each experiment.

5

F. SDS-PAGE of surface expressed ActA.

Surface expressed ActA protein was prepared from late-log phase bacterial cultures grown in LB broth (OD₆₀₀ ~0.7) by resuspending equivalent amounts in SDS-PAGE buffer and boiling for 5 min. which extracts surface-
10 expressed proteins but does not perturb the cell wall. Equivalent amounts were loaded on 7% SDS-PAGE and visualized with Coomassie blue.

G. *Xenopus laevis* cell extract motility assays.

X. laevis egg cytoplasmic extract was prepared as described (Theriot, J. A., J. Rosenblatt, D. A. Portnoy, P. J. Goldschmidt-Clermont, and T. J. Mitchison. 1994. Involvement of profilin in the actin-based motility of *L. monocytogenes* in cells and in cell-free extracts. Cell 76(3):505-517.) and supplemented with tetramethylrhodamine iodoacetamide-labeled actin (Theriot, J. A., and D. C. Fung. 1998. *Listeria monocytogenes*-based assays for actin assembly factors. 20 Methods Enzymol. 298:114-122.). SLCC-5764-derived strains were grown overnight to stationary phase, washed 1×, added to cell extracts and incubated for 45 minutes before microscopic observation.

H. LD₅₀ determinations.

25 Limited LD₅₀ were performed in BALB/c mice as described (Portnoy, D. A., P. S. Jacks, and D. J. Hinrichs. 1988. Role of hemolysin for the intracellular growth of *Listeria monocytogenes*. J. Exp. Med. 167(4):1459-1471.). Animal experiments were performed in the laboratory of Archie Bouwer at Oregon Health Sciences Center, Portland, OR.

30

I. Identification of the PSA attachment site and construction of pPL2.

The PSA attachment site (tRNA^{Arg}-attBB') DNA sequence was obtained through a combination of inverse PCR and genome walking. Inverse PCR was

performed on *Sau*3 AI-digested DP-L4061 DNA (WSLC 1042 lysogenic for PSA) (SEQ ID NO:26 is the sequence of 2,072 bp surrounding WSLC 1042 tRNA^{Arg}-*attBB*) using the divergent primers PL95 (5'-ACATAATCAGTCCAAAGTAGATGC)(SEQ ID NO:16) and PL97 (5'-ACGAATGTAAATATTGAGCGG) (SEQ ID NO:17) that anneal within the PSA *int* gene. The resultant DNA sequence was used to design further oligonucleotides and these were used with the Genome Walker kit (Clontech), per the manufacturer's recommendations. DNA sequence and tRNA analysis was done with using MacVector (Accelrys), DNAsis (Hitachi), BLAST algorithm (2), and tRNAscan-SE (Lowe, T. M., and S. R. Eddy. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 25(5):955-964.).

pPL1 was modified to utilize a different attachment site on the *L. monocytogenes* chromosome by replacing the U153 integrase gene and attachment site in the plasmid. The PSA *int* and *attPP'* were PCR amplified from PSA genomic DNA with primers PL100 (5'-GAAGATCTCCAAAAATAAACAGGTGGTGG) (SEQ ID NO:18) and PL101 (5'-CATGCATGCGTGGAGGGAAAGAAGAACGC) (SEQ ID NO:19) with Vent DNA polymerase, digested with *Bgl* II and *Sph* I, and ligated to pPL1 that had been digested with the same enzymes. The resultant plasmid was designated pPL2 (FIG 1B). The sequence of pPL2 is provided as SEQ ID NO:28.

The DNA sequence of the PSA tRNA^{Arg}-*attBB'* from serotype 1/2 *L. monocytogenes* strains was obtained by a plasmid-trap strategy. DP-L4211 (pPL2 integrated in 10403S) genomic DNA was digested with *Nsi* I and *Nhe* I, which do not cleave in the vector, and ligated under dilute conditions to promote self-ligation. The ligations were transformed into XL1-blue and chloramphenicol resistant colonies were selected. The plasmids obtained were sequenced with the convergent primers PL94 (5'-GGAGGGAAAGAAGAACGC) (SEQ ID NO:20) and PL95 (sequence above) for *attPB'* and *attBP'*, respectively, which flank *attPP'* in the PSA genomic DNA sequence. Further, because of the divergence between the sequences downstream of the tRNA^{Arg} gene among serotypes, a serotype 1/2 specific PCR assay across tRNA^{Arg}-*attBB'* was developed from the 10403S DNA sequence and used to determine the prophage status of various *L.*

monocytogenes strains. Primers PL102 (5'-TATCAGACCTAACCCAAACCTTCC) (SEQ ID NO:21) and PL103 (5'-AATCGCAAAATAAAAATCTTCTCG) (SEQ ID NO:22) specifically amplify a 533 bp PCR product in non-lysogenic serotype 1/2 strains. The primer pair NC16 (5'-GTCAAAACATACGCTCTTATC) (SEQ ID NO:23) and PL95 specifically amplify a 499 bp PCR product in strains that are either lysogenic or contain an integration vector at tRNA^{Arg}-attBB'. (SEQ ID NO:27 provides the 643 bp surrounding 10403S tRNA^{Arg}-attBB')

10 II. RESULTS AND DISCUSSION

A. pPL1 forms stable, single-copy integrants in various *L. monocytogenes* strains.

pPL1 is the first shuttle integration vector that we constructed to facilitate strain construction in *L. monocytogenes*. In order to test the pPL1 vector, we needed a *L. monocytogenes* strain that did not have a phage at the *comK* bacterial attachment site. We adapted historical methods to cure *L. monocytogenes* strains of their prophages and found after superinfection with phage U153, which has the same attachment site as the endogenous 10403S prophage, we were able to isolate prophage-free strains (see Materials and Methods). The prophage-cured 10403S strain was designated DP-L4056 and was used in subsequent experiments.

Conjugation was chosen as a method for introducing the vector into the target cells, as many *Listeria* spp. are inefficiently transformed. Conjugation of pPL1 from *E. coli* into *L. monocytogenes* was successful. Drug resistant transconjugants arose at a reproducible frequency of $\sim 2 \times 10^{-4}$ per donor *E. coli* cell, approximately three-fold lower than conjugation with autonomously replicating plasmids, indicating $\sim 30\%$ integration efficiency for strains receiving the plasmid by conjugation. All chloramphenicol resistant colonies were positive with the PCR assay using primers PL14 and PL61 and negative using a PCR assay across attPP' in pPL1 (PL14 paired with a primer in the RP4 *oriT*) indicating that they were true integrants and that the full genetic complement of plasmid pPL1 had integrated into the *Listeria* chromosome. In addition, this

experiment demonstrated that pPL1 was not maintained as an episomal plasmid and that the vector did not integrate as a concatamer.

We tested the stability of the integrants under non-selective growth conditions. Three integrant strains, DP-L4074 and the merodiploid strains DP-L4076 and DP-L4078 (described in the following sections) were passed in liquid BHI media for 100 generations and plated for single colonies. Ninety-six colonies were then exposed to 0.1 µg/ml chloramphenicol (to induce CAT gene expression) and patched on plates containing 7.5 µg/ml antibiotic. All colonies retained drug resistance. Thirty colonies from each non-selective growth experiment were assayed with the PL14/PL61 PCR assay and all PCR reactions resulted in the 743 bp product, indicating all transconjugants retained the integrated plasmid.

We further addressed whether the integration vector would be generally useful for any *L. monocytogenes* strain with an available attachment site. There have been greater than 320 listeriophages isolated, and many have restricted host ranges. It was unclear whether there was a biological barrier to U153 integrase gene function in host strains that do not support U153 infection. We therefore picked three strains that did not contain a prophage at the *comK* attachment site; two serotype 4b clinical isolates and SLCC-5764, a serotype 1/2a strain that constitutively expresses the known virulence factors in an unregulated manner and has been useful for studying these virulence factors *in vitro*. Each of these strains was first made streptomycin resistant for counter selection in conjugation experiments (as described in Materials and Methods). Streptomycin resistant derivatives were chosen on the basis of having the same growth rate as the parent strain to avoid experimental complications related to viability. The resultant strains, DP-L4088, DP-L4089, and DP-L4082 all proved suitable recipients for pPL1 integration at a similar frequency to DP-L4056.

Finally, we conducted a survey of *L. monocytogenes* isolates to identify suitable strains that do not harbor a prophage at the *comK* attachment site using the PCR assays across *comK-attBB'* (primers PL60/PL61) and the hybrid *attPB'* (primers PL14/PL61). The results of these experiments (Table 2) indicated many of the strains commonly used to study the biology and pathogenesis of *L. monocytogenes* including 10403S, L028, and EGDe had a prophage at *comK*.

Table 2. Prophage status of various strains

Strain	Description	Source	serotype	PL60/PL61 <i>comK</i>	PL14/PL61 <i>attPB'</i>
10403S	wild type	rabbit pellets	1/2a	- ^a	+
DP-L4056	10403S phage cured	This work	1/2a	+ ^b	-
DP-L861	SLCC-5764 (Mack)	WT (overexpresser)	1/2a	+	-
DP-L3818	WSLC 1118::A118	Camembert cheese	4b	-	+
DP-L3633	EGDe	WT (1960s human)	1/2a	-	+
DP-L3293	LO28	WT (clinical origin)	1/2c	-	+
DP-L185	F2397	L.A., Jalisco cheese	4b	+	-
DP-L186	ScottA	Massachusetts outbreak, milk	4b	+	-
DP-L188	ATTC 19113	Denmark, human	3	+	-
DP-L1168	clinical	cole slaw	4b	+	-
DP-L1169	clinical	patient	4b	+	-
DP-L1170	clinical	patient	4b	+	-
DP-L1171	clinical	brie	1/2b	+	-
DP-L1172	clinical	alfalfa tablets	4b	+	-
DP-L1173	clinical	deceased patient	4b	-	+
DP-L1174	clinical	deceased patient	4b	-	+
DP-L3809	1981 Halifax	placenta	4b	+	-
DP-L3810	1981 Halifax	CSF & brain	4b	+	-
DP-L3812	1981 Halifax	coleslaw	4b	+	-
DP-L3813	1996 Halifax	blood	?	-	+
DP-L3814	1981 Halifax	CSF	4b	+	-
DP-L3815	1993 Halifax	CSF	1/2a	+	-
DP-L3816	1995 Halifax	blood	?	+	-
DP-L3817	1993 Halifax	CSF	1/2a	+	-
DP-L3862	1998 Michigan	patient	4b	-	+

^a (-): negative PCR result for primer pair noted at top of column. ^b (+): positive PCR result for primer pair noted at top of column. The PL60/PL61 primer pair specifically amplify a 417 bp PCR product in non-lysogenic strains and result in no PCR product in lysogenic strains. The PL14/PL61 primer pair specifically amplify a 743 bp PCR product in lysogenic strains and result in no PCR product in non-lysogenic strains.

B. The status of *comK* did not affect the virulence of *L. monocytogenes*.

10 We next compared DP-L4056 and DP-L4074 to wild-type 10403S in standard virulence assays to determine if the presence of a prophage at *comK*, lack of prophage, or integration vector altered the virulence phenotypes. These three strains were assayed for LLO activity, ability to form plaques in monolayers of L2 cells, and for virulence in the mouse LD₅₀ assay (Table 3). All were
15 indistinguishable from one another, strongly suggesting that the integrity of the *comK* ORF and the presence of pPL1 had no measurable impact on virulence.

Table 3. Complementation of *actA* and *hly*

Strain	Genotype	Hemolysis on blood plates	Hemolytic activity ^a	Plaque size ^b	LD ₅₀ ^c
10403S	wild type	+	nd	100.(na)	~2x10 ⁴
DP-L4056	10403S phage cured	+	97	101 (1.4)	<1x10 ⁵
DP-L4074	DP-L4056 <i>comK</i> ::pPL1	+	98	99 (1.4)	<1x10 ⁵
DP-L4027	DP-L2161 phage cured, Δhly	-	0	0 (0)	1x10 ⁸
DP-L4075	DP-L4027 Δhly , <i>comK</i> ::pPL24	+	99	97 (3.9)	<1x10 ⁵
DP-L4076	DP-L4056 <i>comK</i> ::pPL24	+	198	96 (2)	nd
DP-L4029	DP-L3078 phage cured, $\Delta actA$	nd	nd	0 (0)	2x10 ⁷
DP-L4077	DP-L4029 $\Delta actA$, <i>comK</i> ::pPL25	nd	nd	86 (4)	<1x10 ⁵
DP-L4078	DP-L4056 <i>comK</i> ::pPL25	nd	nd	72 (6.8)	nd

^a Hemolytic units data shown is from one representative experiment. nd: not

determined. ^b Plaque size is the average of 8 to 10 independent experiments and

5 shown as a percent of wild type (defined as 100%). Standard deviations are

shown in parentheses. na: not applicable. ^c LD₅₀s of 10403S and Δhly (DP-

L2161) were determined in (37), the LD₅₀ of the $\Delta actA$ strain (DP-L1942, a

smaller deletion within the *actA* ORF that does not support actin nucleation at the bacterial surface) was determined in (4).

10

C. Full complementation of *hly* at the phage attachment site.

Listeriolysin-O (LLO), the gene product of *hly*, is a secreted pore-forming cytolysin that is responsible for escape from the membrane-bound vacuole when *L. monocytogenes* first enters a host cell. LLO is absolutely required for the

15 intracellular life cycle of *L. monocytogenes* and virulence. LLO activity can be measured by hemolytic activity on red blood cells. *Hly* mutants fail to form plaques in monolayers of L2 cells and are 5 logs less virulent in the mouse LD₅₀ assay.

We cloned the *hly* structural gene into pPL1 and conjugated this plasmid

20 from *E. coli* to phage-cured wild type and Δhly *L. monocytogenes* derivatives, resulting in DP-L4076 (an *hly* merodiploid) and DP-L4075 (*hly* only at the phage *comK*-att site). These strains were tested for hemolytic activity on blood plates, for the relative amount of hemolytic units secreted, ability to form a plaque in a

monolayer of L2 cells, and virulence in the mouse LD₅₀ assay (Table 3). The quantitative complementation of *hly* in the deletion strain background and the doubling of hemolytic units produced in the merodiploid strain indicate two things. First, gene expression is not *de facto* affected by ectopic expression at the *comK* chromosomal position. Second, the *hly* promoter is self-contained. Additionally, a two-fold increase in the amount of LLO is not deleterious to the virulence and intracellular life cycle of *L. monocytogenes*, at least as measured by plaquing.

D. Complementation of *actA* at the phage attachment site approaches wild-type expression.

ActA, a second major *L. monocytogenes* virulence factor, is responsible for commandeering host cell actin-cytoskeletal factors used for intracellular bacterial motility. ActA is also absolutely required for bacterial pathogenesis as mutants in *actA* are both unable to spread from cell-to-cell and form a plaque in a cell monolayer and are 3 logs less virulent than wild type. Additionally, ActA expression appears to be more complex than that of LLO: there are two promoters that drive *actA* expression. One is immediately upstream of the *actA* ORF and the second is in front of the *mpl* gene upstream of *actA*.

We constructed several strains to evaluate the complementation of *actA* at the phage attachment site. The first group included making second-site complemented (DP-L4077) and merodiploid (DP-L4078) strains in the 10403S background. These were assayed for plaque formation in an L2 monolayer (Table 3). Integrated ActA did not fully complement in this assay (plaque size of 86%) and the merodiploid strain formed an even smaller plaque (72% of wild type). We interpret these results to indicate that there may be a small contribution of the second promoter upstream of the *mpl* gene for optimal *actA* expression. Additionally, there appears to be a critical concentration of ActA on the surface of intracellular bacteria because two copies of *actA* (with 3 promoters driving expression) further decreases the ability to spread from cell to cell, presumably because there is too much ActA on the bacterial surface for optimal motility.

We further tested ActA complementation in the virulence gene over-expressing strain SLCC-5764. ActA is effectively expressed in this strain from the *comK-attBB'* site (Fig 3A, lane 9). Considering the plaquing data of 10403S

complemented *actA* strains, it may have been predicted that the merodiploid strain DP-L4085 would make more ActA than the parent strain. However, this was not observed: the parent strain, the complemented strain and the merodiploid strain all expressed similar levels of ActA (Fig 3A, lanes 5, 8, and 9).

5 This observation was likely due to the complete lack of regulation and high level of constitutive expression of ActA in SLCC-5764. Additionally, DP-L4087 supports actin nucleation at the bacterial surface, actin tail formation and bacterial motility in cell extracts (Fig. 3B). The results of these cell-extract experiments indicate that the integration vector system for complementation will

10 be useful for *in vitro* studies of *L. monocytogenes* motility, facilitating strain construction and placing various molecular constructs in different host strains for study in a desired set of assays. In particular, several alleles of *actA* that have unusual motility phenotypes have been transferred to the SLCC-5764 $\Delta actA$ strain using pPL1 and are currently being evaluated in cell extracts. The study of

15 these mutants in the simplified cell-extract system should yield insights into the activities of poorly understood regions of the ActA protein.

The strains referenced above are provided in the following Table 1.

Strain	Relevant genotype or plasmid
<u><i>E. coli</i>:</u>	
SM10	Conjugation donor. <i>F- thi-1 thr-1 leuB6 recA tonA21 lacY1 supE44 Mu+C I- [RP4-2(Tc::Mu)] Km^r Tra+</i>
XL1-Blue	Plasmid manipulations. <i>recA1 endA1 gyrA96 thi-1 hsdR17 supE44 relA1 lac [F'proAB lacI^qZ ΔM15 Tn10 (Tet^r)]</i>
DP-E4067	Integration vector pPL1/SM10
DP-E4068	<i>hly</i> integration vector pPL24/SM10
DP-E4069	<i>actA</i> integration vector pPL25/SM10
DP-E4190	Integration vector pPL2/SM10
<u><i>L. monocytogenes</i>:</u>	
10403S	wild type
DP-L4056	10403S phage cured
DP-L4027	DP-L2161 phage cured, Δhly
DP-L4029	DP-L3078 phage cured, $\Delta actA$
DP-L4074	DP-L4056 <i>comK</i> ::pPL1
DP-L4075	DP-L4027 Δhly , <i>comK</i> ::pPL24
DP-L4076	DP-L4056 <i>comK</i> ::pPL24
DP-L4077	DP-L4029 $\Delta actA$, <i>comK</i> ::pPL25
DP-L4078	DP-L4056 <i>comK</i> ::pPL25
SLCC-5764	Virulence gene over expresser (Mack, DP-L861)
DP-L862	Mack-4R (SLCC-5764 rough isolate)
DP-L4082	SLCC-5764 Str ^r derivative
DP-L3780	SLCC-5764 $\Delta actA$ (deletion of amino acids 7-633)

Strain	Relevant genotype or plasmid
DP-L4083	DP-L3780S (Str ^r derivative)
DP-L4084	DP-L4082, <i>comK</i> ::pPL1
DP-L4085	DP-L4082, <i>comK</i> ::pPL25
DP-L4086	DP-L4083, <i>comK</i> ::pPL1
DP-L4087	DP-L4083 $\Delta actA$, <i>comK</i> :: <i>actA</i>
DP-L4088	DP-L1169S 4b strain, Str ^r
DP-L4089	DP-L1172S 4b strain, Str ^r
DP-L4090	DP-L4088, <i>comK</i> ::pPL1
DP-L4091	DP-L4089, <i>comK</i> ::pPL1
DP-L4199	EGDe, Str ^r derivative
DP-L4026	WSLC 1042, (ATCC 23074)
DP-L4061	WSLC 1042::PSA
DP-L4221	10403S, tRNA ^{Arg} ::pPL2

E. Phage PSA integrates into a tRNA^{Arg} gene and pPL2 construction.

pPL1 integration into *L. monocytogenes* strains that harbour a prophage in the *comK* attachment site is hindered by the process of first having to cure the

5 prophage from the host strain. To alleviate the need for the phage-curing step, the specificity of pPL1 integration was changed to that of the PSA prophage. PSA, (Phage from ScottA) is the prophage of *L. monocytogenes* strain ScottA, a serotype 4b strain that was isolated during an epidemic of human listeriosis. Using the PSA genomic DNA sequence, we identified an integrase-like ORF with

10 a contiguous non-coding sequence that we predicted to contain the *attPP'* sequences. The PSA integrase sequence was then used to obtain the DNA sequence of PSA-*attBB'* from the PSA lysogenic strain DP-L4061 (see Materials and Methods). PSA was found to integrate in a tRNA^{Arg} gene that is 88% identical to a tRNA^{Arg} gene (*trnSL-ARG2*) from *B. subtilis*. The anticodon of the

15 tRNA^{Arg} gene is 5'UCU, the most commonly used arginine anticodon in *L. monocytogenes*. The PSA and bacterial attachment sites share 17 bp of DNA identity, and the tRNA^{Arg}-*attBB'* contains a short nucleotide sequence that completes the tRNA^{Arg} sequence that is interrupted by integration of PSA (Fig 4). The attachment site tRNA^{Arg} gene is present only once in the genome of *L.*

20 *monocytogenes* strain EGDe and apparently only once in the serotype 4b. This indicates that not only is the PSA integration site unique, but also that precise reconstitution of the gene upon integration (or excision) is likely required for survival of the cell.

pPL2 was constructed by replacing the U153 listeriphage integrase gene and attachment site in pPL1 with the PSA listeriphage integrase gene and attachment site. pPL2 was transformed into SM10 and the resulting strain was mated into 10403S, EGDe (carrying a streptomycin resistance mutation) and the serotype 4b strain DP-L4088. Chloramphenicol resistant transconjugants arose from each of these crosses at approximately 2×10^{-4} per donor cell, the same rate as pPL1 integration. Two recombinants from each cross were restreaked under drug selection and tested by PCR for the presence of PSA-*attBP'* using primers NC16 and PL95. The expected 499 bp PCR product was obtained in each of the colonies tested, indicating pPL2 integrates into tRNA^{Arg}-*attBB'* in both serotype 1/2 and 4b strains. We tested the stability of the integrated pPL2 in both EGDe and DP-L4088 strains with the same non-selective 100-generation experiment described for pPL1. Forty-nine colonies from each of the amplified cultures were tested for chloramphenicol resistance. The EGDe-derived strains retained 100% drug resistant colonies indicating complete stability of the integrants. In the case of the DP-L4088 integrant, two of the 49 colonies were chloramphenicol sensitive, suggesting a low level of excision can occur in this serotype 4b strain. In order to test whether precise excision had occurred, we PCR amplified across tRNA^{Arg}-*attBB'* and sequenced the PCR products. The wild-type DNA sequence was obtained, indicating a precise excision event.

During the course of our PCR experiments, we noted a divergence between the tRNA^{Arg}-*attPB'* sites from serotype 4b and serotype 1/2 *L. monocytogenes*. To determine nature of this divergence, we isolated and sequenced the tRNA^{Arg}-*attBB'* site from 10403S (as described in Materials and Methods). We found that the sequence of *attPB'* in 10403S (3' of the tRNA^{Arg} gene) is unrelated to that of the serotype 4b strain WSLC 1042. In contrast to this, the sequence of *attBP'* (5' of the tRNA^{Arg} gene) in 10403S is 96-97% identical to the corresponding regions in *L. monocytogenes* serotype 4b strain WSLC 1042, the serotype 4b strain sequenced by TIGR, and the serotype 1/2a strain EGDe sequenced by the European *Listeria* Consortium. Thus, the bacterial *attBB'* sequences recognized by the PSA integrase are likely to encompass more of the *attB* DNA sequence than the *attB'* DNA sequence. Additionally, we tested the availability of the tRNA^{Arg}-*attBB'* in the common

laboratory strains of *L. monocytogenes* with a PCR assay using primers PL102 and PL103. We found the tRNA^{Arg} attachment site to be available in strains 10403S, EGEE, and L028 indicating that pPL2 may be readily utilized in these backgrounds for strain construction, complementation, and genetics studies
5 without first curing endogenous prophages.

F. Expression of Aquoria victoria GFP.

A GFP coding sequence as described in United States Patent No. 5,777,079 (the disclosure of which is herein incorporated by reference), is cloned
10 into plasmid pPL1, transferred into the genome of *L. monocytogenes*. The GFP coding sequence is amplified by polymerase chain reaction (PCR) and the PCR fragment is cloned into the multiple cloning site of pPL1. A suitable promoter, containing appropriate transcriptional elements and a translational leader sequence for expressing the GFP in *L. monocytogenes* are cloned at the 5' end
15 of the GFP coding sequence such that they induce the expression of the GFP protein. The modified pPL1 plasmid constructs are electroporated into *E. coli* strain SM10 using standard techniques, and the modified pPL1 plasmid construct is conjugated into *L. monocytogenes* as described above. Recombinant *L. monocytogenes* are selected on BHI plates supplemented with 7.5 µg/ml
20 chloramphenicol and 200 µg/ml streptomycin. Individual colonies are picked and screened by PCR for integration at the phage attachment site using the primers PL14 (5'-CTCATGAACTAGAAAAATGTGG-3') (SEQ ID NO:13), PL60 (5'-TGAAGTAAACCCGCACACGATG-3')(SEQ ID NO:14) and PL61 (5'-TGTAACATGGAGGTTCTGGCAATC-3')(SEQ ID NO:15). Cultures of
25 recombinant *L. monocytogenes* are grown, prepared and screened for GFP.

III. Utility of pPL1 and pPL2.

The construction and characterization of the first single step site specific integration vectors for use in *L. monocytogenes* furthers the genetic tools
30 available for the study of this pathogen. These vectors allow more facile strain construction than historic methods and are widely useful in various strains used to study the intracellular life cycle of *L. monocytogenes*. Additionally, stable merodiploid strains can be constructed to allow refined copy number studies and

studies of interactions within a protein through multimerization and testing of the dominance or recessive nature of different alleles of a gene in the same bacterial strain.

pPL1 and pPL2 are also useful for vaccine development, e.g., for at least
5 enhancing, including both eliciting and boosting, an immune response to a target
cells or cells, e.g., a foreign pathogen, etc. Several recombinant *L.*
monocytogenes systems have been used to elicit cell-mediated immune
responses in mice (Frankel, F. R., S. Hegde, J. Lieberman, and Y. Paterson.
1995. Induction of cell-mediated immune responses to human immunodeficiency
10 virus type 1 Gag protein by using *Listeria monocytogenes* as a live vaccine
vector. J. Immunol. **155**(10):4775-4782.; Goossens, P. L., G. Milon, P. Cossart,
and M. F. Saron. 1995. Attenuated *Listeria monocytogenes* as a live vector for
induction of CD8+ T cells in vivo: a study with the nucleoprotein of the
lymphocytic choriomeningitis virus. Int. Immunol. **7**(5):797-805; Ikonomidis, G., Y.
15 Paterson, F. J. Kos, and D. A. Portnoy. 1994. Delivery of a viral antigen to the
class I processing and presentation pathway by *Listeria monocytogenes*. J. Exp.
Med. **180**(6):2209-2218.; Shen, H., M. K. Slifka, M. Matloubian, E. R. Jensen, R.
Ahmed, and J. F. Miller. 1995. Recombinant *Listeria monocytogenes* as a live
vaccine vehicle for the induction of protective anti-viral cell-mediated immunity.
20 Proc. Natl. Acad. Sci. USA **92**(9):3987-3991.). One limitation with plasmid-based
expression of recombinant proteins in *L. monocytogenes* is the stability of the
plasmids *in vivo* (i.e. in the host animal) without selection. Additionally,
chromosomal construction of strains expressing foreign antigens is time
consuming. pPL1 and pPL2 alleviate both of these concerns.

25

It is evident from the above results and discussion that subject invention
provides a number of advantages. The construction and characterization of the
first single step site specific integration vectors for use in *L. monocytogenes* as
30 described herein furthers the genetic tools available for the study of this
pathogen. For example, the subject vectors and methods allow more facile strain
construction than historic methods and are widely useful in various strains used
to study the intracellular life cycle of *L. monocytogenes*. Furthermore, the subject
invention provides important new tools for the production of vaccine preparations.

One limitation with plasmid-based expression of recombinant proteins in *L. monocytogenes* is the stability of the plasmids *in vivo* (i.e. in the host animal) without selection. Additionally, chromosomal construction of strains expressing foreign antigens is time consuming. The subject vectors and methods of use
5 alleviate both of these concerns. As such, the present invention represents a significant contribution to the art.

All publications and patent application cited in this specification are herein incorporated by reference as if each individual publication or patent application
10 were specifically and individually indicated to be incorporated by reference. The citation of any publication is for its disclosure prior to the filing date and should not be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention.

15 Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it is readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

20

WHAT IS CLAIMED IS:

1. An integration vector capable of site-specific *Listeria* genome integration.
2. The integration vector according to Claim 1, wherein said integration
5 vector is a plasmid.
3. The integration vector according to Claim 2, wherein said integration
vector comprises a bacteriophage integrase gene and a bacteriophage
attachment site.
- 10 4. The integration vector according to Claim 3, wherein said bacteriophage is
a listeriophage.
5. The integration vector according to Claim 3, wherein said attachment site
15 provides for integration at an integration site selected from the group consisting
of: the comK integration site and the tRNA^{Arg} integration site.
6. The integration vector according to Claim 1, wherein said integration
vector further includes a multiple cloning site.
- 20 7. The integration vector according to Claim 6, wherein said integration
vector further includes a coding sequence.
8. The integration vector according to Claim 7, wherein said coding sequence
25 encodes a polypeptide.
9. The integration vector according to Claim 8, wherein said polypeptide is an
antigen.
- 30 10. The integration vector according to Claim 1, wherein said integration
vector is pPL1.
11. The integration vector according to Claim 1, wherein said integration
vector is pPL2.

12. A method of transforming a *Listeria*, said method comprising:
contacting said *Listeria* with an integration vector according to Claim 1
under conditions sufficient for said integration vector to integrate into said
5 *Listeria*'s genome.
13. A *Listeria* transformed with a vector according to Claim 1.
14. A method of eliciting or boosting a cellular immune response to an antigen
10 in a subject, said method comprising:
administering to said subject an effective amount of *Listeria* cells
according to Claim 13.
15. The method according to Claim 14, wherein said *Listeria* cells are
15 attenuated.
16. A vaccine comprising a strain of *Listeria* cells according to Claim 13,
wherein said *Listeria* cells express a heterologous antigen.
- 20 17. The vaccine according to Claim 16, wherein said *Listeria* cells are
attenuated.
18. A recombinant culture of *Listeria* cells according to Claim 13.
- 25 19. The recombinant culture according to Claim 18, wherein said *Listeria* cells
are attenuated.
20. A kit for use in preparing a vector according to Claim 7, said kit
comprising:
30 a vector according to Claim 1; and
at least one nuclease that cuts said vector at said multiple cloning site.
21. The kit according to Claim 20, wherein said kit further comprises a host
cell.

22. A kit for use in preparing a cell according to Claim 13, said kit comprising:
a vector according to Claim 1;
at least one nuclease that cuts said vector at said multiple cloning site; and
5 a *Listeria* cell.

24. A system for preparing a vaccine according to Claim 16, said system
comprising:
a vector according to Claim 1;
10 at least one nuclease that cuts said vector at said multiple cloning site;
a coding sequence for said heterologous antigen;
and
Listeria cells.

15

Figure 1.




Figure 2.




Figure 3.





Figure 4.



B.

352 AGTGCCATTTCCTGATAGCTCAGCTGGATAGAGCAACGGCCTCTAAGCCGTCGGTCGGGGGTTTCAATCCCTCTCAGGACGTAAATAGCTAT
 19228 TTAACATAAAATGTTTGTGGTATTATTGTGGTATATATATCTAAATGGCTTTATATCAGTGCTGTTAATCCCTCTCAGGACGTAAATAGTAA

JC20 Rec'd PCT/PTO 07 SEP 2005

SEQUENCE LISTING

<110> The Regents of the University of California

<120> SITE SPECIFIC LISTERIA INTEGRATION
VECTORS AND METHODS FOR USING THE SAME

<130> BERK-017WO

<150> 10/136,860

<151> 2002-04-30

<160> 28

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 1

ggacgtcatt aaccctcact .aaagg

25

<210> 2

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 2

ggacgtcaat acgactcact atagg

25

<210> 3

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 3

ggacgtcgct atttaacgac cctgc

25

<210> 4

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 4

gagctgcagg agaattacaa cttatatcgt atgggg

36

<210> 5
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 5
gcactgcagc cgcttgccct catctgttac gcc }

<210> 6
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 6
catgcatgcc tctcgccctgt cccctcagtt cag 33

<210> 7
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 7
gtagatctta actttccatg cgagaggag 29

<210> 8
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 8
gggcatgcga taaaaagcaa tctatagaaa aacagg 36

<210> 9
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 9
cctaagcttt cgatcatcat aattctgtc 29

<210> 10
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 10
gggcatgcag atcttttttt cagaaaatcc cagtacg 37

<210> 11
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 11
ggtctagatc aagcacatac ctag 1

<210> 12
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 12
cgggatcctg aagcttgga agcag 25

<210> 13
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 13
ctcatgaact agaaaaatgt gg 22

<210> 14
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 14
tgaagtaaac ccgcacacga tg 22

<210> 15
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 15
tgtaacatgg aggttctggc aatc 24

<210> 16
<211> 24
<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 16

acataatcag tccaaagtag atgc

24

<210> 17

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 17

acgaatgtaa atattgagcg g

1

<210> 18

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 18

gaagatctcc aaaaataaac aggtggtgg

29

<210> 19

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 19

catgcatgcg tggagggaaa gaagaacgc

29

<210> 20

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 20

ggagggaaaag aagaacgc

18

<210> 21

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide

<400> 21

tatcagacct aacccaaacc ttcc

24

<210> 22
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 22
aatcgcaaaa taaaaatctt ctcg

4

<210> 23
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> oligonucleotide

<400> 23
gtcaaaacat acgctcttat c

21

<210> 24
<211> 6101
<212> DNA
<213> Shuttle integration vector pPL1

<220>
<221> misc_feature
<222> 3676
<223> n = A,T,C or G

<400> 24
gacgtcaata cgactcacta tagggcgaat tgggtaccgg gccccccctc gaggtcgacg 60
gtatcgataa gcttgatata gaattcctgc agcccggggg atccactagt tctagagcgg 120
ccgccaccgc ggtggagctc cagcttttgt tccctttagt gaggggttaat gacgtcgcta 180
tttaacgacc ctgccctgaa ccgacgaccg ggtcgaattt gctttcgaat ttctgccatt 240
catccgctta ttatcactta ttcaggcgta gcaccaggcg tttaagggca ccaataactg 300
ccttaaaaaa attacgcccc gccctgccac tcatcgagc actggtgtaa ttcattaagc 360
attctgccga catgggaagcc atcacagacg gcgatgata cctgaatcgc cagcggcatt 420
agcaccttgt cgccttgctg ataataatttg cccatggtga aaacgggggc gaagaagttg 480
tccatatttg ccacgtttta atcaaaactg gtgaaactca cccagggtat ggctgagacg 540
aaaaacatat tctcaataaa cccttttaggg aaataggcca ggttttcacc gtaacacgcc 600
acatcttgcg aatatatgtg tagaaactgc cggaaatcgt cgtggtattc actccagagc 660
gatgaaaacg tttcagtttg ctcatggaaa acggtgtaac aagggtgaac actatcccat 720
atcaccagct caccgtcttt cattgccata cggattccg gatgagcatt catcaggcgg 780
gcaagaatgt gaataaaggc cggataaaaac ttgtgcttat tttcttttac ggtctttaa 840
aaggccgtaa tatccagctg aacggtctgg ttataggtac attgagcaac tgactgaaat 900
gcctcaaaat gttcttttac atgccattgg gatatatcaa cgggtggtata tccagtgtat 960
tttttctcca ttttagcttc cttagctcct gaaaatctcg ataactcaaa aaatacgc 1020
ggtagtgtat ttatttcatt atggtgaaag ttggaacctc ttacgtgccg atcaacgtct 1080
cattttcgcc aaaagttggc ccagggtctc cgggtatcaa cagggaacac aggtatttat 1140
tattctgcga agtgatcttc cgtcacaggt atttattcgg cgcaaagtgc gtcgggtgat 1200
gctgccaact tactgattta gtgtatgat gtgtttttga ggtgctccag tggcttctgt 1260
ttctatcagc tgtccctcct gttcagctac tgacggggtg gtgcgtaacg gcaaaagcac 1320
cgccggacat cagcgctagc ggagtgtata ctggcttact atgttggcac tgatgagggt 1380
gtcagtgaag tgcttcatgt ggcaggagaa aaaaggctgc accggtgcgt cagcagaata 1440
tgtgatacag gatatattcc gcttcctcgc tcactgactc gctacgctcg gtcgttcgac 1500
tgccggcagc ggaaatggct tacgaacggg gcggagattt cctggaagat gccaggaaga 1560
tacttaacag ggaagtgaga gggccgcggc aaagccgttt ttccataggc tccgcccccc 1620
tgacaagcat cacgaaatct gacgctcaaa tcagtgtgtg cgaaaccga caggactata 1680

aagataccag	gcgtttcccc	ctggcggctc	cctcgtgcgc	tctcctgttc	ctgcctttcg	1740
gtttaccggt	gtcattccgc	tgttatggcc	gcgtttgtct	cattccacgc	ctgacactca	1800
gttccgggta	ggcagttcgc	tccaagctgg	actgtatgca	cgaaccccc	gttcagttccg	1860
accgctgcgc	cttatccggt	aactatcgtc	ttgagtccaa	cccggaaaaga	catgcaaaaag	1920
caccactggc	agcagccact	ggtaattgat	ttagaggagt	tagtcttgaa	gtcatgcgcc	1980
ggttaaggct	aaactgaaag	gacaagtttt	ggtgactgcg	ctcctccaag	ccagttacct	2040
cggttcaaaag	agttggtagc	tcagagaacc	ttcgaaaaac	cgccctgcaa	ggcggttttt	2100
tcgttttcag	agcaagagat	tacgcgcaga	ccaaaacgat	ctcaagaaga	tcattcttatt	2160
aatcagataa	aattatttcta	gatttcagtg	caatttatct	cttcaaatgt	agcacctgaa	2220
gtcagcccca	tacgatataa	gttgtaattc	tccgcgcgtt	gccctcatct	gttacgcccg	2280
cggtagccgg	ccagcctcgc	agagcaggat	tcccggttag	caccgccagg	tgcaataaag	2340
ggacagtga	gaaggaaacac	ccgctcgcgg	gtgggcctac	ttcacctatc	ctgcccggct	2400
gacgccgttg	gatacaccaa	ggaaagtcta	cacgaaccct	ttggcaaaat	cctgtatatc	2460
gtgcgaaaaa	ggatggatat	accgaaaaaa	tcgctataat	gaccccgaa	cagggttatg	2520
cagcggaaaa	gcgctgcttc	cctgctgttt	tggtggaatat	ctaccgactg	gaaacaggca	2580
aatgcaggaa	attactgaac	tgaggggaca	ggcgagaggc	atgcgataaa	aagcaatcta	2640
tagaaaaaca	ggttactttt	tatttataat	tttagtttct	cgattcgttt	ccgtccaacg	2700
agagaaaacg	aggaactaaa	caatctaaat	aaacaagcta	ctagagccat	tcaatagtaa	2760
cttgttcacc	gtcaatataa	attttattaa	ttagtgtatt	taaataaagt	tgcttttctc	2820
ggaactctaa	agagtcaaaa	tcaactgttg	ctaaatcagc	taaattttct	tgtatctttt	2880
tatttttctt	caattcttcg	ttagcttcta	tttgtgtctc	ataataatta	atttgagcat	2940
cgatatcagc	catcatagca	tcaagtctcg	aaactctgta	agaaccgctg	atatataaat	3000
caaatagccg	tttctttttt	acgtgttctg	ttttaagttt	ttcatttaag	ctatctaatt	3060
cgtcttcttt	atctacattc	ctagaagcga	aactatagtt	attcacgcga	tcaataatta	3120
attcctcgag	tttgtcagct	ctccaaattt	tatttccaca	tttttctagt	tcatgagtat	3180
gtttgtaagt	cttgcaacta	taatatctat	aatgatattt	ttttccgcgg	gaaacagtat	3240
cttttctcgg	atgacaaaaa	cccaaccac	atttttccaca	cactacaaaa	ttatttagca	3300
acgtgctga	atgtctattc	atatttggat	ttttacccat	gcgagaaaaa	atttcttgaa	3360
ctcgataaaa	ttgttcctct	gaaataatag	gctcatgaac	accttttgta	tgacttttat	3420
ccgcataaga	tacataacca	cagtataaat	cattagttag	ccaattgttg	taactgctat	3480
atgatttcac	tttgaatcct	aattttttta	gtctcttctg	taaagtggta	atgctttttt	3540
cttctcctaaa	aatatcataa	atcatttgta	attgttttgc	ttcttcttca	ttaatatata	3600
atttagtatc	tataacatca	tagccgaatg	ttctaccttt	tgcatgtcgt	aaaggaagac	3660
ctgcttcaat	acgctnaatt	ttcccatca	ccatacgatc	acgtatagtt	tcgcgtctca	3720
attgagcaaa	tacggataat	ataccaatca	tcgcgcgcgc	aaatgggcta	gaggtgtcaa	3780
gagtttcaga	caaactaaca	aattctacat	tgttttttta	gaagtattct	tcaataagcg	3840
ttatcgtatc	tctttgtgag	cgggaaagtc	tatctaagcg	atatacaaca	acagcatcaa	3900
tttcatgtaa	tttacttagc	atttcattta	gtgcggggcg	attcatgttt	gaaccgctgt	3960
atccgccgtc	ttgaaaaata	tcgtatacgt	cccaatcctt	cgagcggcac	aaggctgtta	4020
gcttttcagt	ttgagcttgt	atagagtaat	tctctatttg	ttcttgagta	gatacgcgta	4080
tataaatagc	tgccctcatt	tcggttctcc	tctcgcatgg	aaagttaaga	tctttttttc	4140
agaaaatccc	agtacgtaat	taagtatttg	agaattaatt	ttatattgat	taataactaag	4200
tttaccagct	tttcacctaa	aaaacaaatg	atgagataat	aactccaaag	gctaaagagg	4260
actataccaa	ctatttgtaa	taattctgta	acagttgaaa	agcgaacgtg	tattcttagg	4320
gcttgagatg	tactgctggg	taaacctttt	tagtgtaagt	gggatgtgaa	cgtaatacaa	4380
caactttcgc	tatgggaaac	ctattgtttt	ttgttaatag	aaaaacttaa	tacatttgta	4440
atataaaaac	cggcagtttt	tccgttcttc	gtgactcgaa	atgaattgcc	agatgagttt	4500
atgggtattct	ataatagaag	gtatggagga	tgttatataa	tgagacagaa	ttatgatgat	4560
cgaaagctag	cttggcactg	gccgtcgttt	tacaacgtcg	tgactgggaa	aaccctggcg	4620
ttaccaact	taatcgctt	gcagcacatc	cccccttcgc	cagctggcgt	aatagcgaag	4680
aggcccgac	cgatcgccct	tcccaacagt	tgcgcagcct	gaatggcgaa	tggcgcctga	4740
tcggttatct	tctccttacg	catctgtgcg	gtatttcaca	ccgcataatca	aatgggttcgg	4800
atctggagct	gtaatatataa	aacctttctc	aactaacggg	gcagggttagt	gacattagaa	4860
aaccgactgt	aaaaagtaca	gtcggcatta	tctcatatta	taaaagccag	tcattaggcc	4920
tatctgacaa	ttcctgaata	gagttcataa	acaatcctgc	atgataacca	tcacaaacag	4980
aatgatgtac	ctgtaaagat	agcggtaaat	atattgaatt	acctttatta	atgaattttc	5040
ctgctgtaat	aatgggtaga	aggtaattac	tattattatt	gatatttaag	ttaaacccag	5100
taaatgaagt	ccatggaaata	atagaaagag	aaaaagcatt	ttcaggtata	ggtgtttttg	5160
gaaacaattt	ccccgaacca	ttatatattct	ctacatcaga	aagggtataaa	tcataaaact	5220
ctttgaagtc	attctttaca	ggagtccaaa	taccagagaa	tgtttttagat	acaccatcaa	5280
aaattgtata	aagtggctct	aacttatccc	aataacctaa	ctctccgtcg	ctattgtaac	5340
cagttctaaa	agctgtattt	gagtttatca	cccttgtcac	taagaaaata	aatgcagggt	5400

aaaattttata	tccttcttgt	tttatgtttc	ggtataaaac	actaatatca	atttctgtgg	5460
ttatactaaa	agtcgtttgt	tggttcaaat	aatgattaaa	tatctctttt	ctcttccaat	5520
tgtctaaaatc	aattttatta	aagttcattt	gatatgcctc	ctaaaatttt	atctaaaagt	5580
aatttaggag	gcttacttgt	ctgctttctt	cattagaatc	aatccttttt	taaaagtcaa	5640
tattactgta	acataaatat	atattttaaa	aatatcccac	tttatccaat	tttcgtttgt	5700
tgaactaatg	ggtgcttttag	ttgaagaata	aagaccacat	taaaaaatgt	ggtcttttgt	5760
gttttttttaa	aggattttgag	cgtagcgaaa	aatccttttc	tttcttatct	tgataataag	5820
ggtaactatt	gcccagatcc	gaaccatttg	atatggtgca	ctctcagtac	aatctgctct	5880
gatgccgcat	agttaagcca	gccccgacac	ccgccaacac	ccgctgacgc	gccctgacgg	5940
gcttgctctgc	tcccggcatc	cgcttacaga	caagctgtga	ccgtctccgg	gagctgcatg	6000
tgtcagaggt	tttcaccgtc	atcaccgaaa	cgcgcgagac	gaaagggcct	cgtagacgc	6060
ctatttttat	aggttaatgt	catgataata	atggtttctt	a		6101

<210> 25

<211> 3897

<212> DNA

<213> Bacteriophage U153

<220>

<221> misc_feature

<222> 695

<223> n = A,T,C or G

<400> 25

aagcttttaa	gaaattcaag	aagaaacatc	ggtaactagc	cataaattaa	ccaaagtctt	60
aatctcgctt	gaagagaaca	aactgattga	aaaaattgga	caatctagag	caacaaaaata	120
caaatttaatt	gaatctacag	aggaatatct	aaccaatctt	caacacacat	ttcgaaaaat	180
tgttcaattt	tatgttgaaa	atgataaata	aaaatatgaa	tgttttttta	ttgttttagta	240
gtgtaacttt	ccatgcgaga	ggagaacgga	aatgaaggca	gctattttata	tacgcgtatc	300
tactcaagaa	caaatagaga	attactctat	acaagctcaa	actgaaaagc	taacagcctt	360
gtgccgctcg	aaggattggg	acgtatacga	tattttcata	gacggcggat	acagcggttc	420
aaacatgaat	cgccccgcac	taaatgaaat	gctaagtaaa	ttacatgaaa	ttgatgctgt	480
tgttgtatat	cgcttagata	gactttcccc	ctcacaaaaga	gatacgataa	cgcttattga	540
agaatacttc	ttaaaaaaca	atgtagaatt	tgttagtttg	tctgaaactc	ttgacacctc	600
tagcccattt	gggcgcgcga	tgattggtat	attatccgta	tttgctcaat	tagagcgaga	660
aactatacgt	gatcgatatg	tgatggggaa	aattnagcgt	attgaagcag	gtcttccttt	720
aacgactgca	aaaggtagaa	cattcggcta	tgatgttata	gatactaaat	tatatattaa	780
tgaagaagaa	gcaaaacaat	tacaaatgat	ttatgatatt	tttgagggaag	aaaaaagcat	840
taccacttta	cagaagagac	taaaaaaatt	aggattcaaa	gtgaaatcat	atagcagtta	900
caacaattgg	ctaactaatg	atttatactg	tggttatgta	tcttatgcgg	ataaaagtga	960
tacaaaagggt	gttcattgagc	ctattatttc	agaggaaaca	ttttatcgag	ttcaagaaat	1020
tttttctcgc	atgggtaaaa	atccaaatat	gaatagagat	tcagcatcgt	tgctaaataa	1080
tttggttagtg	tgtggaaaat	gtgggttggg	ttttgttcat	cggagaaaag	atactgtttc	1140
ccgcggaaaa	aaatatcatt	atagatatta	tagttgcaag	acttacaaac	atactcatga	1200
actagaaaaa	tgtggaaaata	aaatttggag	agctgacaaa	ctcgaggaat	taattattga	1260
tcgcgtgaat	aactatagtt	tcgcttctag	gaatgtagat	aaagaagacg	aattagatag	1320
cttaaatgaa	aaacttaaaa	cagaacacgt	aaaaaagaaa	cggctatttg	atttatatat	1380
cagcggttct	tacgaagttt	cagaacttga	tgctatgatg	gctgatatcg	atgctcaaat	1440
taattattat	gaagcacaaa	tagaagctaa	cgaagaattg	aagaaaaata	aaaagataca	1500
agaaaaattta	gctgatttag	caacagttga	tttgactctt	ttagagttcc	gagaaaagca	1560
actttattta	aaatcactaa	ttaataaaat	ttatattgac	ggtgaacaag	ttactattga	1620
atggctctag	tagcttgttt	atttagattg	tttagttcct	cgttttctct	cgttggacgg	1680
aaacgaatcg	agaaaactaaa	attataaata	aaaagtaacc	tgtttttcta	tagattgctt	1740
tttatcaatt	atatagaaga	aagccgcttt	ttattagatt	ataattgatg	ttttttgatt	1800
tatattttcac	tccctgtgca	aataatgata	taacagcaac	ctcgaacttt	ttagttcggg	1860
gtattttttt	gaaattaatt	tataaaaaaca	cttgcaatta	tataatacat	gtattataat	1920
ataaatatag	aaaggagtgt	agaaagtga	agacatctta	gaggaaataa	aaacagtcct	1980
tgaaattgta	actcttgacg	tagcgtgtat	aacattacgc	aagatagaca	aaaacaagga	2040
caagtaacca	gaggggtgaa	actccccctc	ctctataaaa	gtatatcacg	tctttcataa	2100
attatgaata	aatatatctg	ggttatatta	attgttatat	gcgttaacgg	actcgctagt	2160
tactttcaga	acacagcatt	gaccatcatt	gctatactga	ctacatttagc	ttgttttagta	2220
tatttaataa	aaaataggaa	gtgattaatt	atgacgaaaa	aaacgacctc	tgacgcgcag	2280
ttgaaagcaa	ataaggaatg	gcaaagcaag	aacaaagaac	atgcaaaacta	tttaaaatct	2340

```

cggttcagctg cgcgttcctt tataaagaat aaagctacgt tggaagattt gaaggaactt 2400
gaaaaattaa ttatagaggg aaaaattaat cataaggga tgattaagga taaatgatgc 2460
acgctaagca catgcttggc gttttttgca taaaaaaagc cctaacgttg aagttagga 2520
ctgacatata taaaaaatag aagttgacaa cttaaaggcg actaccacga caggcagctt 2580
acaagctatg actagccttg actaatcatt tatgcgacac tcaaagaatt attatctaac 2640
ttcttaatca agaataacaa aaatcaaaca agttagcaag tatttcaggc attttattta 2700
taacaaatat ctagatcaca aaaatgtcgc ggaaaaaat ggtcacaacc aatattacat 2760
aaacttaaaa gtctcttatt tctcttatca ggtttatgtg ctgttacgtg atttctacat 2820
actctaaaaa ctgtattagc gaataagtct acaacttgaa ttaaatcttt attttgtaga 2880
tccttatatg atgtttcaac agaagagaaa attggatgtt ccattgtaaa tttaatagtt 2940
aaatattctt gtaagctatt taatgattca attgcggtat ttctatcatc tatttgcatt 3000
ttcaaatagt tatttgctgg gttaattggt atttttagaaa ttcatcttac cgttagataa 3060
ataaaataat taaaagacaa agatgtatta ttcaaaagat gattgactag ttggtggtta 3120
tcgactatct taaaatgaaa tttagcatc gattttggtg aaagcatatt aaatattaat 3180
tttttcattt caaaaggcat ctccgaacct ttatctctt tgtaatatc taacttacta 3240
gatggatacc ttttaagata ttttaatttt gcatctctga actgtctaata tacattatat 3300
ggtttctctg tttctaaaaa agcaataaca aaatatctgt tattaaaatt tttattttta 3360
gttatagttc ctgattcatc taaaaaaagt ctcacccag ttctccact tttttactta 3420
aattatatta tactaattaa gtttgaggaa gtggaacgta tgtacttata attcgaagtt 3480
atgaaaaatc cccccatcaa tataaaacaa aaaagcccc gaaataataa tcgagggcat 3540
taaaactaaat ctttttaaca aacttcggtg tttagcagtga gatagtaacc agatttcggt 3600
ttcaagcgag gtgttcgcc ttttgttttc gccattcctg taatcgtgaa gatagtgcct 3660
accggatatg tgccaccggt tttatgcttc tcagtaaagt ctactgaatt gtatagatca 3720
cactgtacta gtgttttaac ttttcgcgga ttttctgtgt agtatgtgtt tttgcttgct 3780
ggtgtgtgtg gttttcctgc ttttaacttc gctaataatg ttgtgttctg cgttgctggt 3840
cctttataat ccttaattcc gtattgattt gctagttttt tacgattcgc aaagctt 3897

```

<210> 26

<211> 2702

<212> DNA

<213> *Listeria monocytogenes*

<400> 26

```

gatatcgcg acgtgaatta aacgcagatt ttgccttttt ttggtcaccg catgaactag 60
gagtagacat gctagacgac accatcattt taaacccagg aagcatttcc ttaccaagag 120
gacgcatccg tgtcaaaaca tacgctctta tcgattcaac accagaaggc attcaagttc 180
gattcatgga cggggacgac aacgaactaa cggacctaac ccaaaccttc ccattaacga 240
agcataacta ggtcaaaaga caccgaaaa agaaaaaat caataactta aagaaaacca 300
ttgacaaaca agcgatttaa acataaaatg gtatttggtt gttgaaaaaa cagtgccatt 360
tgtcctgata gctcagctgg atagagcaac ggccttctaa gccgtcggtc gggggttcga 420
atccctctca ggacgtaaat agctatatta aagaaatctc taaaacgttg aaaaaccttg 480
atattaaagg ttggatggat gttttagaga tttttttata tcttataata tctgttttat 540
tccgtatttt tcatgacatt tgtgacaaaa tttgtgctat ttccatccat ttttaatgtg 600
aaaaaagcat ctatttttagt ttgattatgt tgatgcaaat tagagcttag attattataa 660
tattttaatg ttattaatat caggttgacc tctcctaagt gttagacatg tttcaccagt 720
ctccatagga gtgtggtagc tgattgcaca gtaattatat actttacgtc aatatcaaaa 780
gcaagtccaa ttaaaatgga ttaccttgcc ccgtaaatga caacttctga aaataggtta 840
aaggaacaaa agatgatgta attagggtct agtgcatttg tgggtgaattt aggttttgat 900
tataatgaga atctccgttt agaggttggt cttttgaaaa cgatagaagc aattataggt 960
atcgactacc atatattact gaaaaaagag ctagatttaa taaaaaaata attctaacat 1020
cataggaggc aattatgact tttttaaaca ccttaaaatt aaatttgtaa aatgaaaaaa 1080
agagaatggt atccgatgct tttatgaaaa aacaagaagg aatcattgta aactatatag 1140
tgacttgcag taaggattct gctattggca ttagtaaaaa ggcaattgat atattattga 1200
taatcaatga aaatacattt cctgaatggc caaatgtaga tagatggctt tctattttgc 1260
caaaatattt tacggattct ttttcaaaat caaaaatatt gcatagtga gattggctat 1320
ttgaagagtg gttatactgg tttgaacctg aaaatagatt ttggttttta ggagaattag 1380
atcctgttga taatgagcat ttgaaaataa gcatagtgtt acaagaacac ccttttccag 1440
tagaactcatt agaagttcta cttatgaagc taggaacaag cgaattacat gaaattggtg 1500
tggaatgagg ttaaatgtac ttttaacgga tatatctttt acaatagagc tgaattttgt 1560
tagagtttaa aatgaaaaaa caactaagtt ataacgaaag gagctaacac ttgatggaaa 1620
attacgtgtc aatagtaaaa atcgaaaaca atcttccgt gtgcttttac aacagctcgg 1680
agaaagtagt agcaattgct aagaaaatga atgagattaa cgaagaagct tatatgcatg 1740

```



```

gttacaattg ggaagcattt ttcaactact atttacctaa atatgctcca gatgtcttag 1800
aaggaaatggg ctctgatccg gaagcgggaa tgtatgtggc gtattacacg ctatcacctg 1860
aaactgaggg acgagcagaa aaacttggtc aagtaattac gaatctcatc gaaaatgaag 1920
aactacttta tcaaataatt gaaaatgaag gcaataatat tagttgggat aattaatcct 1980
ttttctaaaa aatccttatac tatttattcg tatagtatta gcaagagggtg aagaacctgt 2040
ataatataat tgacgatatt ttaaagcatt agatcctatt ggcagatgct cttaaaacgt 2100
taaacagtaa aataaaaaaa ctctaaaaca tttgaaaccc tttgtaatta aaagggtgaat 2160
gttttagaga tttttttatc ttgcatttcc catttttatt ccgttggttt tgtggcaaat 2220
tttattaaaa ctagtccaag taattacgaa tctcattgaa aacgaagaac tactttataa 2280
aatagtcaaa aattaggaca agcagattat tgagatgatt gatcctttac tttaataata 2340
atTTTTatgt aaactcatcc cttattagggt gttctattgt atgacttgag agtagttttt 2400
ttgagaattt caagcaataa atttaaatat attagagagt ctaaaattag cactaatccc 2460
taaaaagata tgaacgatat gtgaacgatg ataccaagaa atgaaaaaat ttctatacta 2520
tattcaaatt gtaacgttgg gactgctata attgactctt attgaggcga tataatgcca 2580
catacattaa atacagaata aactcattct ttaagataat aattacatct aaggagacta 2640
atcatgaaaa gaaagataag ttctatcatt gtagtcggga taatgttctt tcaatcatta 2700
ac 2702

```

<210> 27

<211> 643

<212> DNA

<213> *Listeria monocytogenes*

<400> 27

```

agcatttcct taccaagagg ggcgatccgt atcaaaacat acggctctta tcaattcaca 60
ccagaaggca tccaagttcg attcatggac cgagatgaca acgaactatc agacctaac 120
caaaccttcc cattaacgaa taactagggtc taactagggtc aaaagacacc cgaaaaagaa 180
aaaatgcaat aacttaaga aaaccattga caaacaagcg atttaaacat aaaatggtat 240
ttggctgttg aaaagacagt gccatttgtc ctgatagctc agctggatag agcaacggcc 300
ttctaagccg tcggctcgggg gtccgaatcc ctctcaggac gtaatatgaa gcgccgtaaa 360
cgttgttaat acaatgttta cggcgctttt tggtttttcg aagttcaaat aaagtacaaa 420
aaatttaaat tccattaatc tttttcatta attatatgta attaggcttc taaagtcatt 480
actatagtgt tttggcccaa tcttaatttt gaagaatata atctttaatt ttggtattag 540
tcttatttag tagcatttgc tccataaaaa caatagaaaa attaatacca gtcttatata 600
aaaatcttct catgacgaga agatttttat tttgcgattg agc 643

```

<210> 28

<211> 6123

<212> DNA

<213> Shuttle integration vector pPL2

<400> 28

```

gacgtcaata cgactcacta tagggcgaat tgggtaccgg gccccccctc gaggtcgacg 60
gtatcgataa gcttgatata gaattcctgc agcccggggg atccactagt tctagagcgg 120
ccgccaccgc ggtggagctc cagcttttgt tccctttagt gagggttaat gacgtcgcta 180
tttaacgacc ctggccctgaa ccgacgacc ggtcgaaatt gctttcgaat ttctgccatt 240
catccgctta ttatcactta ttcaggcgta gcaccaggcg tttaagggca ccaataactg 300
ccttaaaaaa attacgcccc gccctgccac tcatcgagc actggtgtaa ttcattaagc 360
attctgccga catggaagcc atcacagacg gcatgatgaa cctgaatcgc cagcggcatc 420
agcaccttgt cgccttgctg ataataattt gccatggtga aaacgggggc gaagaagttg 480
tccatatttg ccacgtttta atcaaaactg gtgaaactca cccagggtat ggctgagacg 540
aaaaacatat tctcaataaa cccttttagg aaataggcca ggttttcacc gtaacacgcc 600
acatcttgcg aatatatgtg tagaaactgc cggaaatcgt cgtggtattc actccagagc 660
gatgaaaacg tttcagtttg ctcatggaaa acggtgtaac aaggggtgaa actatcccat 720
atcaccagct caccgtcttt cattgccata cggaaattccg gatgagcatt catcaggcgg 780
gcaagaatgt gaataaaggc cggataaaac ttgtgcttat ttttctttac ggtcttttaa 840
aaggccgtaa ttccagctg aacggctctg ttataggtag attgagcaac tgactgaaat 900
gcctcaaaat gttctttacg atgccattgg gatatatcaa cgggtgtata tccagtgtat 960
tttttctcca ttttagcttc cttagctcct gaaaatctcg ataactcaa aaatacgccc 1020
ggtagtgatc ttatttcatt atgggtgaaag ttggaacctc ttacgtgccg atcaacgtct 1080
cattttcgcc aaaagttggc ccagggtctc ccggtatcaa cagggaacac aggattttat 1140
tattctgcga agtgatcttc cgtcacaggt atttattcgg cgcaaagtgc gtcgggtgat 1200

```

```

gctgccaact tactgattta gtgtatgatg gtgtttttga ggtgctccag tggcttctgt 1260
ttctatcagc tgtccctcct gttcagctac tgacgggggtg gtgcgtaacg gcaaaagcac 1320
cgccggacat cagcgctagc ggagtgtata ctggcttact atgttggcac tgatgagggg 1380
gtcagtgaag tgcttcatgt ggcaggagaa aaagggtgc accggtgcgt cagcagaata 1440
tgtgatacag gatataattcc gcttcctcgc tcaactgactc gctacgctcg gtcgttcgac 1500
tgcggcgagc ggaaatggct tacgaacggg gcggagattt cctggaagat gccaggaaga 1560
tacttaacag ggaagtgaga gggccgcggc aaagccgttt ttccataggc tccgcccccc 1620
tgacaagcat cacgaaatct gacgctcaaa tcagtggtgg cgaaaccgga caggactata 1680
aagataccag gcgtttcccc ctggcggtc cctcgtgcgc tctcctgttc ctgcctttcg 1740
gtttaccggt gtcattccgc tgttatggcc gcgtttgtct cattccacgc ctgacactca 1800
gttccgggta ggcagttcgc tccaagctgg actgatatga cgaaccccc gttcagtccg 1860
accgctgcgc cttatccggt aactatcgct ttgagtccaa cccggaaga catgcaaaag 1920
caccactggc agcagccact ggtaattgat ttagaggagt tagtcttgaa gtcattgcgc 1980
ggttaaggct aaactgaaag gacaagtttt ggtgactgcg ctccctcaaag ccagttacct 2040
cggttcaaaag agttggtagc tcagagaacc ttcgaaaaac cgccctgcaa ggcgggtttt 2100
tcgttttcag agcaagagat tacgcgcaga ccaaaacgat ctcaagaaga tcatcttatt 2160
aatcagataa aatatttcta gatttcagtg caatttatct cttcaaatgt agcacctgaa 2220
gtcagcccca tacgatataa gttgtaattc tccgcgcgtt gccctcatct gttacgcccg 2280
cggtagccgg ccagcctcgc agagcaggat tcccgttgag caccgccagg tgcaataaag 2340
ggacagtga gaaggaaacac ccgctcgcgg gtgggcctac ttcacctatc ctgcccggct 2400
gacgccgttg gatacccaa ggaaagtcta cacgaaccct ttggcaaat cctgtatatc 2460
gtgcgaaaaa ggtggatat tccgaaaaaa tcgctataat gacccgaag cagggttatg 2520
cagcggaaaa gcgctgttcc cctgctgttt tgtggaatat ctaccgactg gaaacaggca 2580
aatgcaggaa attactgaac tgaggggaca ggcgagaggc atgctgtggag ggaaagaaga 2640
acgctgttga aaaaaattc tctggactac ttgaaacaaa agaattaaag tcattttata 2700
aaaaccttga gaaaaacat cttgatataa aaactattta taacgaatat ttatttcaat 2760
gtaataataa ataataattt ttattacata aaatgtttgt ggtattattt gtggtatata 2820
tattcctaat ggctttatat cagtgtgtgt taatccctct caggacgtta aatagtaatg 2880
taaagaaatc tctaaaacgt tgaaaagcct tgatatataa gggcggtatg atgttttgga 2940
gtttttttta tatcgtataa taccggtttt attccgttgt tgaaaaaagc atctactttg gactgattat 3060
aatttgtggt attttcatct gtttttagtg tgaaaaaagc tggtgtatta atgtcatcat 3120
gttgtcttaa attagagctt agatgactat agtattttta ttctacacat aagcctgtat 3180
gaccaagcct atcagctaca taaataatat ccatacccg cttcagaatt gattgtacta catatcttct 3240
gcgtatgtcg tagcttgtgt aatgtcactg gttcagaatt gttgtaagt atgaataata 3300
tcaaagcttt attacaagac gcgttgtcta ctggttatt gtggttaagt agtaataata 3360
acatcaatgg attcttaata gcatgttcct tcatataatc agtatgcaa tttaaatacg 3420
aatgtaataa ttgagcggta gagttatcaa tatagatcac tcgtgatttt tttgttttgg 3480
tatcaatgaa tgtattagtg tacttgtaat cccaagcttt attcacagtt attgaacgtt 3540
tagtgaattt aatatccttc tttgttagtg caataatttc ttogaacctc atgcctgtct 3600
ggacagctag aaagataact gctcgtgata tagaatgaaa ttttgcaagt tcttctaata 3660
gtaaatgaa tttgtctgtt tccataaatt gtgctttatt tttcgctacg tcctgtccgc 3720
ttatatgagc ccttatagtg gggtttttct ctacagtgga tattgcatag tctacagata 3780
aaatcgctct aattttgcgg tgtctggtgt catcaatcga attagttta attttttcat 3840
aatgattaat aaattgttga tattgaaccg gatcgataaa attaatagta gattgactac 3900
cgaaataatc aacgaattga ttataagcaa gcgtataaaa ttctttgaag ttccattctt 3960
ttttcccatc tttaaatgtt ttcattgaata ttaataattt agatgcttta tacattaagt 4020
tcagagaact actatcatgc tgaactgttt ctttccattc accatcgact tttatacgta 4080
ttgtttcact tgtattctgc aaacgctttt tttttatctt cattaatacc accacctgtt 4140
ggcgaacaca atattttaccg tttgctaatt tttttaccta ccagtacgta attagttatt tgagaattaa 4200
tatttttgga gatctttttt tcagaaaatc gttttcacct aaaaaacaaa tgatgagata 4260
ttttatattg attaatacta agtttaccca aactatttgt aataattctg taacagttga 4320
ataactcaa aggctaaaga ggactatacc tgtactgctg ggtaaacctt tatagtgtaa 4380
aaagcgaacg tgtattctta aacaactttc gctatgggaa acctattgtt ttttgtaata 4440
gtgggagtgt aacgttaatc taatataaaa accggcagtt tttccgttct tcgtactcg 4500
agaaaaactt aatacatttg ttatggtatt ctataataga aggtatggag gatgttatat 4560
aatgaattg ccagatgagt atcgaaagct agcttggcac tggccgtcgt tttacaacgt 4620
cgtgactggg aaaaccctgg cgttacccaa cttaatcgcc ttgcagcaca tccccctttc 4680
gccagctggc gtaatagcga agaggcccg accgatcgcc cttcccaaca gttgcgcagc 4740
ctgaatggcg aatggcgct gatgcggtat tttctcctta cgcatctgtg cggtatttca 4800
caccgcatat caaatggttc ggatctggag ctgtaataa aaaaacctct tcaactaacg 4860
gggcaggtta gtgacattag aaaaccgact gtaaaaagta cagtcggcat tatctcatat 4920

```

tataaaagcc	agtcattagg	cctatctgac	aattcctgaa	tagagttcat	aaacaatcct	4980
gcatgataac	catcacaaac	agaatgatgt	acctgtaaag	atagcggtaa	atatattgaa	5040
ttacctttat	taatgaatth	tcctgctgta	ataatgggta	gaaggtaatt	actattatta	5100
ttgatattta	agttaaaccc	agtaaatgaa	gtccatggaa	taatagaaag	agaaaaagca	5160
ttttcaggta	taggtgtttt	gggaaacaat	ttccccgaac	cattatattt	ctctacatca	5220
gaaaggata	aatcataaaa	ctctttgaag	tcattcttta	caggagtcca	aataccagag	5280
aatgttttag	atacaccatc	aaaaattgta	taaagtggct	ctaacttatc	ccaataacct	5340
aactctccgt	cgctattgta	accagttcta	aaagctgtat	ttgagtttat	cacccttgte	5400
actaagaaaa	taaatgcagg	gtaaaattta	tatccttctt	gttttatgtt	tcggtataaa	5460
acactaatat	caatttctgt	ggttataacta	aaagtcgttt	gttggttcaa	ataatgatta	5520
aatatctctt	ttctcttcca	attgtctaaa	tcaattttat	taaagtthcat	ttgatatgcc	5580
tcctaaatth	ttatctaaag	tgaatttagg	aggcttactt	gtctgctttc	ttcattagaa	5640
tcaatcctth	tttaaaagtc	aatattactg	taacataaat	atatatttta	aaaatatccc	5700
actttatcca	attttcgttt	gttgaactaa	tgggtgcttt	agttgaagaa	taaagaccac	5760
attaaaaaat	gtggctcttt	gtgttttttt	aaaggatttg	agcgtagcga	aaaatcctth	5820
tctttcttat	cttgataata	agggtacta	ttgccagat	ccgaaccatt	tgatatgggtg	5880
cactctcagt	acaatctgct	ctgatgccgc	atagttaagc	cagccccgac	acccgccaac	5940
acccgctgac	gcgccctgac	gggcttgtct	gtccccggca	tccgcttaca	gacaagctgt	6000
gaccgtctcc	gggagctgca	tgtgtcagag	gttttcaccg	tcatcaccga	aacgcgcgag	6060
acgaaagggc	ctcgtgatac	gcctatthtt	ataggthaat	gtcatgataa	taatggthtc	6120
tta						6123